RARE BRYOPHYTES AND LICHENS IN IDAHO: STATUS OF OUR KNOWLEDGE

by

Robert K. Moseley and Andrew Pitner Conservation Data Center

February 1996

Idaho Department of Fish and Game 600 South Walnut, P.O. Box 25 Boise, Idaho 83707 Jerry M. Conley, Director

Idaho Bureau of Land Management Idaho Department of Fish and Game

Purchase Order No. D910P50087

TABLE OF CONTENTS

TABLE OF CONTENTS I
LIST OF TABLES I
INTRODUCTION 1
BRYOPHYTES 2
LICHENS 11
CONCLUSIONS AND RECOMMENDATIONS
BRYOPHYTE AND LICHEN REFERENCES AT THE CDC

LIST OF TABLES

Table 1. Status and distribution of rare mosses currently on the Idaho Native PlantSociety rare plant list.	2
Table 2. Additional rare Idaho mosses from Christy and Harpel's Columbia River Basin analysis.	3
Table 3. Status and distribution of rare lichens in Idaho. Image: Distribution of the state of the	11

INTRODUCTION

The compilation and dissemination of information regarding rare cryptogams in Idaho has lagged far behind that for vascular plants. The Conservation Data Center (CDC) has been actively researching sources of information on rare mosses, liverworts, and lichens for several years, including an extensive, computerized literature search for rare species done in 1992 (methods described in Moseley *et al.* 1992. Idaho Rare Vascular Flora: Bibliography, 1896-1991. GTR-INT-292. USDA, Forest Service, Ogden, UT). Because of the large workload associated with vascular plants we were never able to computerize the rare bryophyte and lichen populations into our rare species occurrence data base, or the numerous references into the bibliographic data base. To rectify this, the Idaho BLM provided funding in 1995 to put rare bryophytes and lichens on par with vascular species.

This report represents a summary of the information we compiled. It is organized into four main sections. The first presents the status of our knowledge of rare Idaho bryophytes (mosses and liverworts). It includes a current list of rare species in the state, with conservation ranks and county distributions, and a printout from the CDC data base for all reported Idaho occurrences of some of the rare species. The second section, covering rare lichens, is similar to the bryophyte section. A short Conclusions and Recommendations section follows these two. Finally, we present output from our bibliographic data base of all references pertaining to bryophytes and lichens that we have on file. These citations represent a broad range of material, including sources specific to Idaho's rare species, general articles on nonvascular plant conservation, air quality monitoring, ecological studies, and others.

BRYOPHYTES

In 1989, Joe Elliot of Helena, Montana, helped put a preliminary list of rare Idaho bryophytes together. These ten moss species have been on the Idaho rare plant list ever since (Table 1). Because few field inventories have been conducted for mosses, their conservation status has changed little since they were added to the list. Only in the last couple of years have any occurrences been submitted to the CDC by field biologists.

Table 1. Status and distribution of rare mosses currently on the Idaho Native Plant Society rare plant list. Conservation Ranks and County Distribution abbreviations are defined in the "blue book" (Conservation Data Center. 1994. Rare, threatened, and endangered plants and animals in Idaho. Idaho Department of Fish and Game, Boise). Under Comments, EO refers to the number of occurrences in the CDC data base, which appear at the end of the Bryophyte section.

Species	Rank	Distribution	Comments
Bryum calobryoides	G3/S1	El	1 EO
Buxbaumia aphylla	G4/S?	Northern Idaho	
Buxbaumia piperi	G4/S3	Id, La	10 EOs, 3 not in db
Hellodium blandowii	G5/S2	Bn, Cu, Id, Lm	4 EOs, 1 not in db
Hookeria lucens	G4/S1	Cw, Id	5 EOs
Meesia longiseta	G5/S?	Northern Idaho	
Sphagnum mendocinum	G4/S1	Bn, Id	2 EOs
Tayloria serrata	G5/S3?	Northern Idaho	
Ulota megalospora	G4/S2?	Northern Idaho	
SUGGESTED TO BE DROPPE	D:		

Ulota curvifolia - no documented occurrences in Idaho!?

David Wagner from Eugene, Oregon, recently brought to our attention the presence of a globally rare liverwort in Idaho, *Sphaerocarpos hians*. Two historical collections from the Palouse Prairie near Pullman, Washington, and Moscow, Idaho, were all that were thought to exist until a collection was made recently in the Willamette Valley of western Oregon (John Christy, Oregon Natural Heritage Program, personal communication, 1996). More inventory needs to be conducted, but for the time being we've ranked it G1/SH, with one historical collection in Idaho from Latah County.

As part of the Interior Columbia River Basin assessment, John Christy and Judith Harpel (1995.

Bryophytes of the Columbia River Basin, south of the Canadian border) were contracted by federal agencies to compile a list of bryophytes occurring in the study area and make recommendations on the conservation of rare species. Although there was some overlap with the current Idaho list, their recently completed study suggested 31 additional moss species that should be of conservation concern in Idaho (Table 2). Most species are known from only one or two collections. Of special note is the narrow endemic, *Orthotrichum holzingeri*, known only from Nez Perce County, Idaho, and adjacent Washington.

pecies	Distribution	Number of Collections
acolia menziesii var. baueri	Ko	1
chodontium olypicum	Sh	1
lymodon tophaceus	Id	1
epanocladus fluitans	Ko, La	2
alypta intermedia	Cu	2
ssidens ventricosus	Bn	1
matocaulis vernicosus	Bn, Cu, Ko	3
nalothecium arenarium	El	1
grohypnum alpinum	Bn	1
rohypnum duriusculum	Id	?
grohypnum styriacum	Cu	?
gotrichum aligerum	Sh	1
hotrichum flowersii	Ow	?
hotrichum hallii	Cu	1
hotrichum holzingeri	Ne	1
hotrichum pulchellum	Ne	1
notrichum striatum	Id, Ko	2

Table 2. Additional rare Idaho mosses from Christy and Harpel's Columbia River Basin analysis.

Table 2. Continued

Species	Distribution	Number of Collections	
Plagiothecium cavifolium	Sh	1	
Plagiothecium undulatum	La	1	
Pogonatum contortum	Sh	1	
Polytrichum formosum	Bd	1	
Pottia lanceolata	Cu	1	
Pottia nevadensis	Go	1	
Pseudoleskea radicosa var. denudata	Bd	1	
Pseudoleskea saviana	Va	1	
Racomitrium aquaticum	Id	1	
Racomitrium varium	Ко	1	
Schistidium cinclidodonteum	Ко	1	
Splachnum sphaericum	La	1	
Tayloria acuminata	E	1	
Triptercladum leucocladulum	Ne	1	

Following are abbreviated occurrence records for the five mosses listed in Table 1 that have been entered into the CDC data base and the single *Sphaerocarpos hians* occurrence. Several collections have been difficult to locate and are not yet in the data base. We are beginning to compile information on the species listed in Table 2 and will be processing occurrence records for these species during 1996.

SPHAEROCARPOS HIANS

Occurrence Number: 001 Survey Site Name: PARADISE CREEK County: Latah USGS quadrangle: MOSCOW WEST First Observed: CA 1928 Last Observed: CA 1928 Population Data: Ca 1928: Collected in Latah County by Clark. Habitat Description: On soil. Specimens: Clark s.n. (WS).

BRYUM CALOBRYOIDES BEAUTIFUL BRYUM

Occurrence Number: 001 Survey Site Name: CHATTANOOGA HOT SPRINGS County: Elmore USGS quadrangle: ATLANTA EAST Last Observed: 1941-07-05 First Observed: 1941-07-05 Population Data: 1941: No data. Collected by F. A. MacFadden. Habitat Description: Above hot springs. Specimens: F. A. MacFadden 18542 (UBC) - determined by A. L. Andrews.

BUXBAUMIA PIPERI

PIPER'S BUG-ON-A-STICK Occurrence Number: 001 Survey Site Name: EAST OF SOB POINT County: Idaho USGS quadrangle: STILLMAN POINT First Observed: 1994-07-21 Last Observed: 1994-07-21 Population Data: 1994: 3 mature genets in fruit. Population vigor assessed as average. Area thoroughly surveyed by Leonard Lake, Nez Perce NF. Habitat Description: Old growth Thuja plicata stand. Thuja plicata/Athyrium filix-femina habitat type. Found on downed logs in advanced stages of decay. Specimens:

Occurrence Number: 002 Survey Site Name: FALLS CREEK County: Idaho USGS quadrangle: STILLMAN POINT First Observed: 1994-08-02 Last Observed: 1994-08-02 Population Data: 1994: 2 mature genets in fruit. Population vigor assessed as average. 1 log thoroughly surveyed by Leonard Lake, Nez Perce NF. Habitat Description: Old growth Thuja plicata stand. Thuja plicata/Adiantum pedatum habitat type. On downed log in advanced stages of decay. Specimens:

Occurrence Number: 003 Survey Site Name: BIG SMITH CREEK - LITTLE SMITH CREEK County: Idaho USGS quadrangle: SYRINGA First Observed: 1994-08-16 Last Observed: 1994-08-19 Population Data: 1994: 2 populations: 14 or more individuals in fruit near Little Smith Creek; 49 or more in fruit near Big Smith Creek. Observations by Karen Gray, Clearwater NF, Lochsa RD. Habitat Description: Southerly aspects. Slopes 25-35%. Partial to shaded light. Moist slumpy substrate. Growing on rotten Thuja plicata logs. Thuja plicata/Asarum caudatum/Asarum caudatum habitat type. Associated species include Equisetum telmateia, Athyrium filix-femina, Adiantum pedatum, Rubus ursinus, Polystichum munitum, Asarum caudatum, Smilacina stellata, Clintonia uniflora, Montia cordifolia, Festuca subuliflora, Carex hendersonii, Cornus canadensis, Dryopteris expansa, Kindbergia oregana, and Rhytidiopsis robusta. Specimens: Karen L. Gray 319 (ID). Karen L. Gray 320 (ID, MRC).

Occurrence Number: 004 Survey Site Name: NORTH OF BURGDORF HOT SPRINGS County: Idaho USGS quadrangle: BURGDORF First Observed: 1991-06-05 Last Observed: 1991-06-05 Population Data: 1991: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: On a Pinus contorta log. Potential forest climax is Abies lasiocarpa/Vaccinium scoparium habitat type. Specimens: Roger Rosentreter 6865 (SRP). Occurrence Number: 005 Survey Site Name: FOUR-BIT CREEK County: Idaho USGS quadrangle: MUSSELSHELL First Observed: 1995-08-06 Last Observed: 1995-08-06 Population Data: 1995: 20 individuals or more in fruit. Observation by Karen Gray. Habitat Description: Aspect: west. Substrate: rotten log. Shady, moist Thuja plicata forest. Specimens: Karen L. Gray 868 (MRC).

Occurrence Number: 006 Survey Site Name: CAMP CREEK County: Idaho USGS quadrangle: GOODWIN MEADOWS First Observed: 1959-07-16 Last Observed: 1959-07-16 Population Data: 1959: Collected by Higinbotham. Habitat Description: Abies grandis/Pachistima association; on decaying wood. Specimens: Higinbotham 771 (WSU).

Occurrence Number: 007 Survey Site Name: DEVOTO GROVE County: Idaho USGS quadrangle: ROCKY POINT First Observed: 1975-10-18 Last Observed: 1975-10-18 Population Data: 1975: Collected by Hunter and Wagner. Habitat Description: On downed Thuja plicata log. Specimens: Hunter s.n. (WSU). Wagner s.n. (WSU).

Occurrence Number: 008 Survey Site Name: ELDORADO/DOLLAR CREEK County: Idaho USGS quadrangle: MUSSELSHELL First Observed: 1994-08-22 Last Observed: 1994-08-22 Population Data: 1994: Collected by Karen Gray, Clearwater NF. Habitat Description: E aspect in Thuja plicata-Abies grandis forest on a rotten log. Specimens: Karen L. Gray 333a (ID).

Occurrence Number: 009 Survey Site Name: ELDORADO/MAY CREEK County: Idaho USGS quadrangle: MUSSELSHELL First Observed: 1994-09-14 Last Observed: 1994-09-14 Population Data: 1994: Collected by Karen Gray, Clearwater NF. Habitat Description: NE aspect in Thuja plicata-Abies grandis forest on a rotten log. Specimens: Karen L. Gray 384 (ID).

Occurrence Number: 010 Survey Site Name: YAKUS CREEK/STRAY CREEK County: Idaho USGS quadrangle: SYRINGA First Observed: 1994-09-06 Last Observed: 1994-09-06 Population Data: 1994: Collected by Karen Gray, Clearwater NF. Habitat Description: E aspect in Thuja plicata-Abies grandis forest on a rotten log. Specimens: Karen L. Gray 371 (ID).

HELODIUM BLANDOWII

BLANDOW'S HELODIUM Occurrence Number: 001 Survey Site Name: JACK CREEK County: Idaho USGS quadrangle: TRAPPER CREEK First Observed: 1994-05-26 Last Observed: 1994-05-26 Population Data: 1994: Ca 500 mature vegetative genets. Population vigor assessed as good. Observation on a cursory visit by Leonard Lake, Nez Perce NF. Habitat Description: Minerotrophic seep with sphagnum under Pinus contorta. Rolling Pinus contorta landform adjacent to Jack Creek riparian area. Specimens:

Occurrence Number: 002 Survey Site Name: BEAR VALLEY CREEK County: Lemhi USGS quadrangle: LEM PEAK First Observed: 1973 Last Observed: 1973 Population Data: 1973: No data. Occurred in macroplot but not any microplots so no cover estimate. Observation by Alma Hanson. Habitat Description: Aspect: 100 degrees. 1% slope. Aulacomnium palustre-Brachythecium velutinum moss association

which occurs on cool moist terraces supporting Abies lasiocarpa/Calamagrostis canadensis habitat type. Associated with Aulacomnium palustre, Brachythecium velutinum, B. hylotapetum, Pohlia nutans, and Hypnum revolutum. Specimens:

Occurrence Number: 003 Survey Site Name: MAYS CREEK FEN County: Custer USGS quadrangle: OBSIDIAN First Observed: 1993-08 Last Observed: 1993-08 Population Data: 1993: Plants rare, comprising <1% of cover in microplot RM002. Area thoroughly surveyed by Bob Moseley and Rob Bursik, Idaho CDC. Habitat Description: 3% slope. Peat depth 55 cm. Pinus contorta/Vaccinium occidentale community type. Near top of high Sphagnum hummocks around the base of Vaccinium occidentale, Betula glandulosa, and Ledum glandulosum. Additional associates include Lonicera caerulea, Potentilla fruticosa, Salix planifolia, S. wolfii, Calamagrostis canadensis, Carex aquatilis, C. deweyana, C. livida, C. muricata, C. rostrata, Deschampsia cespitosa, Eleocharis pauciflora, Muhlenbergia filiformis, Aster foliaceus, Senecio cymbalarioides, and Swertia perennis. Other bryophytes include Aulocomnium palustre, Calliergon stramineum, and Homalothecium nitens. Specimens: Rob Bursik collection number unknown (ID).

Occurrence Number: 004 Survey Site Name: REDFISH LAKE County: Custer USGS quadrangle: MT. CRAMER First Observed: 1972-06-27 Last Observed: 1972-06-27 Population Data: 1972: Collected by Anderegg. Habitat Description: No information available. Specimens: Anderegg M-71 (ID).

HOOKERIA LUCENS

LIGHT HOOKERIA Occurrence Number: 001 Survey Site Name: ELDORADO CREEK/SNOW CREEK County: Idaho USGS quadrangle: MUSSELSHELL First Observed: 1994-09-26 Last Observed: 1994-09-26

Population Data: 1994: 6 square inch vegetative mat. Population vigor assessed as fair to poor. Area thoroughly surveyed by Karen Gray, Clearwater NF, Lochsa RD.

Habitat Description: Occurs on the edge of a seep in Thuja plicata/Abies grandis forest with Glyceria striata, Senecio triangularis, Dryopteris expansa, Boykinia major, Conocephalum conicum, other liverworts, Plagiomnium sp., Rhizomnium sp., Carex amplifolia, Veratrum viride, Polystichum munitum, and Hylocomium splendens. Overall site quality is fair. Specimens:

Occurrence Number: 002 Survey Site Name: YAKUS CREEK/STRAY CREEK County: Idaho USGS quadrangle: SYRINGA First Observed: 1994-09-06 Last Observed: 1994-09-06 Population Data: 1994: 1 square foot vegetative patch observed by Karen Gray, Clearwater NF, Lochsa RD. Habitat Description: In a seepy area in Thuja plicata/Abies grandis forest with Boykinia major, Senecio triangularis, and other mosses and liverworts. East aspect. Overall site quality is good. Specimens: Karen L. Gray 374 (ID, MRC).

Occurrence Number: 003 Survey Site Name: WEITAS CREEK - NW OF JOHNNY BUTTE County: Clearwater USGS quadrangle: LEAN-TO POINT First Observed: 1994-08-26 Last Observed: 1994-08-26 Population Data: 1994: 3 vegetative patches observed by Karen Gray, Clearwater NF, Lochsa RD. Habitat Description: Found growing near seeps or small creeks on soil or rotten logs. Scattered large-diameter Thuja plicata with Abies grandis (60 year old?). Associates include Alnus rubra, Streptopus amplexifolius, Adiantum pedatum, Gymnocarpium dryopteris, Athyrium filix-femina, and liverworts. Overall site quality is good. Specimens: Karen L. Gray 355, 360, 363 (ID, MRC).

Occurrence Number: 004 Survey Site Name: PETE KING CREEK N County: Idaho USGS quadrangle: SYRINGA First Observed: 1994-08-19 Last Observed: 1994-08-19 Population Data: 1994: Collected by Karen Gray, Clearwater NF, Lochsa RD. Habitat Description: In Thuja plicata-Abies grandis forest. On wet, rotten log at waters edge. SE aspect. Specimens: Karen L. Gray 328 (ID).

Occurrence Number: 005 Survey Site Name: FOUR-BIT CREEK RNA County: Idaho USGS quadrangle: MUSSELSHELL First Observed: 1995-12-01 Last Observed: 1995-12-01 Population Data: 1995: Population is vegetative and assessed as vigorous. Observation on a cursory visit by Karen Gray. Habitat Description: N aspect. Growing on the edge of a small tributary in a seepy, nearly flat area adjacent to water. Thuja plicata, Abies grandis, Abies lasiocarpa, and Picea engelmannii forest. Specimens: Karen L. Gray 1076 (MRC, GFC).

SPHAGNUM MENDOCINUM

PEATMOSS Occurrence Number: 001 Survey Site Name: MOSQIUTO BAY FEN County: Bonner USGS quadrangle: PRIEST LAKE NE First Observed: 1901-08 Last Observed: 1901-08 Population Data: 1901: No data. Collected by C. V. Piper. Habitat Description: Peatland habitat. Specimens: C. V. Piper 349 (WTU, NY).

Occurrence Number: 002 Survey Site Name: SAWYER RIDGE County: Idaho USGS quadrangle: SAWYER RIDGE First Observed: 1974 Last Observed: 1974 Population Data: 1974: No data. Collected by E. F. Layser. Habitat Description: Montane bog community. Perennially wet seepage area. Complex of Abies lasiocarpa (Abla) series

habitat types (Abla-Caltha biflora, Abla-Menziesia ferruginea, Abla-Xerophyllum tenax, and Abla-Luzula hitchcockii). Representative species include Picea engelmannii, Sphagnum spp., Caltha biflora, Dodecatheon jefferyii, Gentiana affins, Kalmia polifolia, Ledum glandulosum, and Tofieldia palustris.

Specimens: E. F. Layser 2216, 2219, 2222, 2224 (WTU).

LICHENS

At the same time that the bryophyte list was being compiled, the preliminary list of rare Idaho lichens was compiled by Roger Rosentreter and Bruce McCune. In contrast to the bryophytes, the list of rare lichens has been more dynamic due to the greater inventory and collecting activity of lichenologists and other biologists in the state. Table 3 presents the current rare lichen list and reflects several changes made as a result of new data received in 1995, including some additions, suggested deletions, and up-to-date conservation ranks.

Table 3. Status and distribution of rare lichens in Idaho. Species not on the current Idaho Native Plant Society list are marked with an asterisk. As in Table 1, Conservation Ranks and County Distribution abbreviations are defined in the "blue book." Under Comments, EO refers to the number of occurrences in the CDC data base, which appear at the end of the Lichens section.

Species	Rank	Distribution	Comments
Aspicilia fruticulosa	G?/S3?	Bk, Cu, Lm, Ow	numerous collections from Idaho at SRP
*Catapyrenium congestum	G2/S1	Aa, Bi, Jf, Ow	3 EOs; 1 not in db
*Cetraria subalpina	G3/S1	Bn, Bd	?
*Cladina rangiferina	G5?/S1	Bn	1 not in db
*Cladonia andreggii	G1/S1	La	1 EO
*Cladonia imbricaria	G?/S1?	Bn, Bw, Id	3 EOs
Cladonia luteoalba	G2/S1	Cu	1 EO
Cladonia transcendens	G5/S2	Cw, Id, La, Lw, Ko, Sh	1 EO; 3 not in db
Cladonia uncialis	G5/S1	Bn, Id, Sh	1 EO; 2 not in db
Collema curtisporum	G1/S1	Id, Ko, Sh	2 not in db
Collema furfuraceum	G5/S1	Sh?	
*Dermatocarpon lorenzianum	G1/S1	Wa	1 EO
Lobaria hallii	G4/S3	Bn, Bw, Ko, Sh	10 EOs; 5 not in db
Lobaria linita	G4/S1	Id	1 EO
Lobaria scrobiculata	G4/S1	Id	3 EOs
Physcia semipinnata	G5/S?	Id?	

Table 3. Continued

Species	Rank	Distribution	Comments	
Pilophorus acicularia	G4/S1	Cw, Sh	2 EOs; 1 not in db	
Pseudocyphellaria anthraspis	G4/S3	Cw, Id	9 EOs; 9 not in db	
Sphaerophorus tuckermannii	G4/S1	Cw, Id?, La?	1 not in db	
Texosporium sancti-jacobi	G2/S2	Aa, El	13 EOs	
Thamnolia vermicularis	G4/S1	Cu	1 EO	
Xanthoparmelia idahoensis	G1/S1	Lm	5 EOs	
SUGGESTED TO BE DROPPED:				
Cladonia verruculosa - more comme	on than previou	usly thought		
Ramalina thrausta - more common than previously thought				
Thelomma ocellatum - more common than previously thought				
Hypogymnia apinnata - more comm	on than previo	usly thought		
Hypogymnia enteromorpha - more c	common than p	reviously thought		

Following are abbreviated occurrence records for the 18 lichens listed in Table 3 that have been entered into the CDC data base (EOs). The status of several common species are being evaluated and some of the numerous known occurrences have not been entered yet. Also, several hard-to-locate specimens are not entered.

CATAPYRENIUM CONGESTUM

Occurrence Number: 001 Survey Site Name: DRY CREEK County: Ada USGS quadrangle: BOISE NORTH First Observed: 1993-04-17 Last Observed: 1993-04-17 Population Data: 1993: Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Found on sandstone rock in Artemisia grasslands. Associated species is Gleocapsa phycobiont. Specimens: Roger Rosentreter 7892 (SRP).

Occurrence Number: 002 Survey Site Name: CIRCULAR BUTTE County: Jefferson USGS quadrangle: CIRCULAR BUTTE First Observed: 1987-06-26 Population Data: 1987: Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Found on deep loamy soils in Eurotia lanata habitat. Specimens: Roger Rosentreter 3950 (SRP) - determined by McCune, 1994. Occurrence Number: 003 Survey Site Name: OREANA SW County: Owyhee USGS quadrangle: OREANA First Observed: 1981-10-01 Last Observed: 1981-10-01 Population Data: 1981: Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Found on soil in Artemisia tridentata habitat. Specimens: Roger Rosentreter 2470 (SRP).

CLADONIA ANDEREGGII

Occurrence Number: 001 Survey Site Name: MOOSE CREEK DIVIDE County: Latah USGS quadrangle: SAND MOUNTAIN First Observed: 1989-06 Last Observed: 1989-06 Population Data: 1989: No data. Collected by Samuel Hammer. Habitat Description: Growing with aquatic bryophytes in deep shade, damp area near stream and seep. Specimens: Samuel Hammer 3798 (FH, paratype). Samuel Hammer 3800 (FH, holotype).

CLADONIA IMBRICARICA

Occurrence Number: 001 Survey Site Name: EMIDA County: Benewah USGS quadrangle: EMIDA First Observed: CA 1989 Last Observed: CA 1989 Population Data: 1989: No data. Collected by Samuel Hammer. Habitat Description: On acidic soil over volcanically-derived rocks. Basalt outcrop. Specimens: Samuel Hammer 3842 (FH) - collection number noted as 2842 in Hammer's 1995 "A synopsis of the genus Cladonia."

Occurrence Number: 002 Survey Site Name: HANNA FLATS CEDAR GROVE BOTANICAL AREA County: Bonner First Observed: CA 1989 Last Observed: CA 1989 Population Data: 1989: No data. Collected by Samuel Hammer. Habitat Description: On acidic soil over volcanically-derived rocks. Specimens: Samuel Hammer 3965 (FH) - collection number noted as 2965 in Hammer's 1995 "A synopsis of the genus Cladonia."

Occurrence Number: 003 Survey Site Name: HARPSTER County: Idaho USGS quadrangle: HARPSTER First Observed: CA 1989 Last Observed: CA 1989 Population Data: 1989: No data. Collected by Samuel Hammer. Habitat Description: On acidic soil over volcanically-derived rocks. Specimens: Samuel Hammer 5311 (FH).

CLADONIA LUTEOALBA

REINDEER LICHEN Occurrence Number: 001 Survey Site Name: ACROSS FROM HOSPITAL BAR County: Lemhi USGS quadrangle: RAMEY HILL First Observed: 1987-07-22 Last Observed: 1987-07-22 Population Data: 1987: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Moss-covered, N-facing small cliff band. Specimens: Roger Rosentreter 4526 (H, OSU).

CLADONIA TRANSCENDENS

TRANSCENDING REINDEER LICHEN Occurrence Number: 001 Survey Site Name: MOOSE CREEK DIVIDE County: Latah USGS quadrangle: SAND MOUNTAIN First Observed: 1989-06 Last Observed: 1989-06 Population Data: 1989: No data. Collected by Samuel Hammer. Habitat Description: On rotting, ecorticate wood, rarely terricolous. Specimens: Samuel Hammer collection number and herbarium unknown.

CLADONIA UNCIALIS

Occurrence Number: 001 Survey Site Name: NEAR CLARK FORK County: Bonner USGS quadrangle: CLARK FORK First Observed: 1981-07-17 Last Observed: 1981-07-17 Population Data: 1981: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: On talus on a N-facing slope. Specimens: Roger Rosentreter 2357 (SRP) - confirmed by Hammer.

DERMATOCARPON LORENZIANUM

Occurrence Number: 001 Survey Site Name: BUCKWHEAT FLATS County: Washington USGS quadrangle: MIDVALE HILL First Observed: 1984-08-27 Last Observed: 1991-06-22 Population Data: 1984 and 1991: Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Found on rock in heavy clay soils, shallow stony basalt is adjacent; Eriogonum thymoides habitat. Specimens: Roger Rosentreter 6896 (SRP). Roger Rosentreter 3538 (SRP).

HYPOGYMNIA APINNATA

TUBE LICHEN Occurrence Number: 001 Survey Site Name: GLOVER CAMPGROUND County: Idaho USGS quadrangle: SELWAY FALLS First Observed: 1974-08-22 Last Observed: 1974-08-22 Population Data: 1974: Collected by Anderegg and Schroeder. Habitat Description: Growing on Cornus sp. in coniferous forest of Thuja plicata, Taxus brevifolia, and Alnus rubra. Specimens: Anderegg and Schroeder 3100 (ID).

Occurrence Number: 002 Survey Site Name: CLEARWATER RIVER/LODGE CREEK County: Idaho USGS quadrangle: SYRINGA First Observed: 1971-07-06 Last Observed: 1971-07-06 Population Data: 1971: Collected by Schroeder. Habitat Description: No information available. Specimens: Schroeder L1697 (ID).

Occurrence Number: 003 Survey Site Name: MERRY CREEK County: Shoshone USGS quadrangle: MERRY CREEK First Observed: 1972-07-15 Last Observed: 1972-07-15 Population Data: 1972: Collected by Doyle Anderegg. Habitat Description: Coniferous forest; Thuja plicata, Pseudotsuga menziesii, and Larix sp. Specimens: Doyle E. Anderegg 722 (ID).

Occurrence Number: 004 Survey Site Name: HEMLOCK BUTTE EAST County: Clearwater USGS quadrangle: ANTHONY PEAK First Observed: 1968-06-09 Last Observed: 1968-06-09 Population Data: 1968: Collected by Anderegg. Habitat Description: No information available. Specimens: Anderegg 101 (ID).

Occurrence Number: 005 Survey Site Name: BANKS GULCH County: Latah USGS quadrangle: SAND MOUNTAIN First Observed: 1969-12 Last Observed: 1969-12 Population Data: 1969: Collected by Anderegg. Habitat Description: No information available. Specimens: Anderegg 112 (ID).

Occurrence Number: 006 Survey Site Name: MANNERING CREEK W County: Latah USGS quadrangle: WEST DENNIS First Observed: 1972-05-11 Last Observed: 1972-05-11 Population Data: 1972: Collected by Anderegg. Habitat Description: Mature conifer forest: Pseudotsuga menziesii, Pinus monticola, and Thuja plicata. Specimens: Anderegg 391 (ID).

Occurrence Number: 007 Survey Site Name: MANNERING CREEK E County: Latah USGS quadrangle: WEST DENNIS First Observed: 1974-05-18 Last Observed: 1974-05-18 Population Data: 1974: Collected by Anderegg. Habitat Description: Mature forest of Thuja plicata and Pseudotsuga menziesii. Specimens: Anderegg 2389 (ID).

Occurrence Number: 008 Survey Site Name: GISBORNE MOUNTAIN County: Bonner USGS quadrangle: PRATER MTN. First Observed: 1969-07-15 Last Observed: 1969-07-15 Population Data: 1969: Collected by Esslinger. Habitat Description: On bark of Pinus monticola, north aspect; Tsuga heterophylla-Pachistima myrsinites habitat type.

Specimens: Esslinger 1393 (ID).

Occurrence Number: 009 Survey Site Name: BOTTLE LAKE RESEARCH NATURAL AREA County: Bonner USGS quadrangle: PRIEST LAKE NE, PRIEST LAKE NW First Observed: 1995-08-15 Last Observed: 1995-08-15 Population Data: 1995: Observation by B. McCune, R. Rosentreter, BLM SO, and A. Debolt, Lower Snake River District BLM.

LOBARIA HALLII

HALL'S LUNGWORT Occurrence Number: 001 Survey Site Name: WEST OF MURRAY County: Shoshone USGS quadrangle: MURRAY First Observed: 1987-08-19 Last Observed: 1987-08-19 Population Data: 1987: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Found on Populus trichocarpa bark. Specimens: Roger Rosentreter 5015 (SRP).

Occurrence Number: 002 Survey Site Name: BEAVER CREEK/TRAIL CREEK County: Shoshone USGS quadrangle: OSBURN First Observed: 1987-08-19 Last Observed: 1987-08-19 Population Data: 1987: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: On Populus trichocarpa bark. Specimens: Roger Rosentreter 5023 (SRP).

Occurrence Number: 003 Survey Site Name: MURRAY County: Shoshone USGS quadrangle: MURRAY, BURKE First Observed: 1985-06-07 Last Observed: 1985-06-07 Population Data: 1985: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: On rock in Thuja forest. Specimens: Roger Rosentreter 3600 (SRP) - determined by C. Wetmare 4/87.

Occurrence Number: 004 Survey Site Name: GROUSE MOUNTAIN County: Bonner First Observed: 1971-06-13 Last Observed: 1971-06-13 Population Data: 1971: Collected by Schroeder. Habitat Description: In shade on a Betula tree in a conifer forest. Specimens: Schroeder L1666 (ID).

Occurrence Number: 005 Survey Site Name: ST. JOE RIVER MILE 45 County: Shoshone USGS quadrangle: CALDER First Observed: 1971-05-09 Last Observed: 1971-05-09 Population Data: 1971: Collected by Schroeder. Habitat Description: On trunk of Populus trichocarpa. Specimens: Schroeder L1548 (ID). Occurrence Number: 006 Survey Site Name: SANTA CREEK County: Benewah USGS quadrangle: SANTA First Observed: 1969-09-13 Last Observed: 1969-09-13 Population Data: 1969: Collected by Schroeder. Habitat Description: On Populus tree in wooded pasture. Specimens: Schroeder L465 (ID).

Occurrence Number: 007 Survey Site Name: ST. MARIES S County: Benewah USGS quadrangle: ST. MARIES First Observed: 1971-04-28 Last Observed: 1971-04-28 Population Data: 1971: Collected by Schroeder. Habitat Description: On trunk of Populus trichocarpa. Specimens: Schroeder L1519 (ID).

Occurrence Number: 008 Survey Site Name: CHATCOLET LAKE County: Benewah USGS quadrangle: BENEWAH LAKE First Observed: 1970-04-18 Last Observed: 1971-03-30 Population Data: 1970 and 1971: Collected by Schroeder and Anderegg. Habitat Description: On Populus tree in open picnic area. Specimens: Schroeder L954 (ID). Schroeder L749 (ID). Anderegg 177 (ID).

Occurrence Number: 009 Survey Site Name: DISTILLERY BAY County: Bonner USGS quadrangle: PRIEST LAKE NW First Observed: 1971-08-02 Last Observed: 1971-08-02 Population Data: 1971: Collected by Schroeder. Habitat Description: On alder tree, in shade. Specimens: Schroeder L1702 (ID).

Occurrence Number: 010 Survey Site Name: ESE OF OROFINO County: Clearwater USGS quadrangle: OROFINO EAST First Observed: 1995-12-03 Last Observed: 1995-12-03 Population Data: 1995: Ca 50 vegetative plants. Population vigor assessed as fair. Observation by Karen Gray. Habitat Description: In wetland on N aspect of ridge. Substrates: Amelanchier alnifolia and Physocarpus malvaceus. Wetland surrounded by Pseudotsuga menziesii and Abies grandis forest. Wetland vegetation: Carex spp., Cornus stolonifera, Salix sp., Acer glabrum, Physocarpus malvaceus, and Amelanchier alnifolia. Associated lichens: Lobaria pulmonaria and Pseudocyphellaria anomola. Specimens: Karen L. Gray 1073, 1109 (MRC, ID).

LOBARIA LINITA SMEARED LUNGWORT Occurrence Number: 001 Survey Site Name: GRAVE PEAK County: Idaho USGS quadrangle: GRAVE PEAK First Observed: 1981-07-24 Last Observed: 1981-07-24 Population Data: 1981: No data. Collected by Bruce McCune. Habitat Description: On alpine sod, ledges on NW side of summit. Specimens: Bruce McCune 11491 (WIS, US, SRP). Bruce McCune 11503 (herbarium unknown).

LOBARIA SCROBICULATA

PORED LUNGWORT Occurrence Number: 001 Survey Site Name: BARTH HOT SPRINGS County: Idaho USGS quadrangle: SHEEP HILL First Observed: 1987-07-29 Last Observed: 1989-08 Population Data: 1987-1989: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: On granitic rock and Betula bark above the river banks. Specimens: Roger Rosentreter 4527 (SRP) - collected 7/29/87. Roger Rosentreter 4915 (SRP) - collected 6/18/88. Roger Rosentreter 4956 (SRP) - collected 7/28/88. Roger Rosentreter 1906 (SRP) - collected 8/89.

Occurrence Number: 002 Survey Site Name: FIVEMILE BAR County: Idaho USGS quadrangle: FIVEMILE BAR First Observed: 1993-08-16 Last Observed: 1993-08-16 Population Data: 1993: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: N-facing cliffs. Pseudotsuga menziesii habitat. Specimens: Roger Rosentreter 7987 (SRP).

Occurrence Number: 003 Survey Site Name: SALMON RIVER MILE 110 County: Idaho USGS quadrangle: CAREY DOME First Observed: 1975-06-28 Last Observed: 1975-06-28 Population Data: 1975: Collected by Anderegg and Schroeder. Habitat Description: On a shaded, N aspect, talus and rubble slope. Specimens: Anderegg and Schroeder 3219 (ID).

PILOPHORUS ACICULARIS

NAIL LICHEN Occurrence Number: 001 Survey Site Name: MURRAY County: Shoshone USGS quadrangle: MURRAY, BURKE First Observed: 1985-06-07 Last Observed: 1985-06-07 Population Data: 1985: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: On rock in Thuja forest. Specimens: Roger Rosentreter 3599 (SRP) - determined by JWT 3/87.

Occurrence Number: 002 Survey Site Name: BEAVER CREEK/NORTH FORK CLEARWATER RIVER CONFLUENCE County: Clearwater USGS quadrangle: SHEEP MTN. First Observed: 1995-07-15 Last Observed: 1995-07-15 Population Data: 1995: Collected by Karen Gray. Habitat Description: On rock near stream. Moist, shady area. Growing with liverworts. Specimens: Karen L. Gray 752 (ID, MRC).

PSEUDOCYPHELLARIA ANTHRASPIS

WHITE-DOT LICHEN
Occurrence Number: 001
Survey Site Name: MOUTH OF APGAR CREEK
County: Idaho
USGS quadrangle: LOWELL
First Observed: 1994-01-12 Last Observed: 1994-03-11
Population Data: 1994: Appeared to have fallen to the ground. Collected by Karen gray, Clearwater NF, Lochsa RD.
Habitat Description: Shady Thuja plicata forest. East aspect. Moist.
Specimens: Karen L. Gray 123B (MRC). Karen L. Gray 157 (MRC).

Occurrence Number: 002 Survey Site Name: MOUTH OF RYE PATCH CREEK County: Idaho USGS quadrangle: LOWELL First Observed: 1994-04-24 Last Observed: 1994-04-24 Population Data: 1994: Collected by Karen Gray, Clearwater NF, Lochsa RD. Habitat Description: East aspect. On branch of hardwood, probably either Alnus, Betula, or Acer. Thuja plicata/Abies grandis forest. With Lobaria pulmonaria. Specimens: Karen L. Gray 182 (ID, MRC).

Occurrence Number: 003 Survey Site Name: WEITAS CREEK - WNW OF JOHNNY BUTTE County: Clearwater USGS quadrangle: LEAN-TO POINT First Observed: 1994-08-26 Last Observed: 1994-08-26 Population Data: 1994: Population vigor assessed as fair. Observation on a cursory visit by Karen Gray, Clearwater NF, Lochsa RD. Habitat Description: NW aspect. On a Thuja plicata branch. Overall site quality is fair. Specimens: Karen L. Gray 364 (herbarium unknown).

Occurrence Number: 004 Survey Site Name: WASHINGTON CREEK County: Clearwater USGS quadrangle: CLARKE MTN. First Observed: 1966-08-09 Last Observed: 1966-08-09 Population Data: 1966: No data. Collected by Peter F. Stickney. Habitat Description: Young Abies grandis forest. Substrate: dead limb of Thuja plicata. Specimens: Peter F. Stickney 1590 (MRC) - determined to be Stricta anthraspis by John W. Thompson in 1968.

Occurrence Number: 005 Survey Site Name: AQUARIUS County: Clearwater USGS quadrangle: THOMPSON POINT First Observed: 1989-07-17 Last Observed: 1995-06-15 Population Data: 1989: No data. Collection by Ann DeBolt. 1995: In fruit. Observation by Sarah Walker, Clearwater NF, and Juanita Lichthardt, Idaho CDC. Habitat Description: E aspect. On fallen Thuja branch in old growth Thuja plicata stand. With Alnus rubra, Boykinia major, Polystichum munitum, and Cornus canadensis. Other lichens present include Sphaerophorus globosus and Lobaria pulmonaria. Specimens: Ann DeBolt 1257 (SRP) - collection label gives elevation as 2000 ft.

Occurrence Number: 006 Survey Site Name: TWO SHADOWS CREEK County: Idaho USGS quadrangle: SYRINGA First Observed: 1979-03-14 Last Observed: 1995-06-06

Population Data: 1979: No data. Collected by Roger Rosentreter, Idaho State BLM. 1995: 4 or 5 thalli were found on the ground. More probably exist in the trees. Observation on a cursory visit by Karen Gray, Clearwater NF, Lochsa RD. Habitat Description: Thuja plicata/Abies grandis forest. River terrace, gentle slope and aspect. Substrate: Thuja branch. Associated with Lobaria pulmonaria. Associated understory species include Cardamine constancei, Carex hendersonii, Kindbergia oregana, Festuca subuliflora, and Eburophyton austiniae. Specimens: Roger Rosentreter 1174 (SRP).Karen L. Gray 705 (MRC). Karen L. Gray 708 (ID).

Occurrence Number: 007 Survey Site Name: ISABELLA LANDING County: Clearwater USGS quadrangle: THOMPSON POINT, SHEEP MTN. First Observed: 1995-06-22 Last Observed: 1995-06-22 Population Data: 1995: Found frequently, 20% vegetative and 80% in fruit. Area surveyed by Sarah Walker, Clearwater NF, and Juanita Lichthardt, Idaho CDC. Habitat Description: Moist Thuja plicata and Abies grandis forest. Found frequently through elevation range. Predominantly on shrub branches like Philadelphus lewisii, one occurrence on a mossy rock. Usually with Peltigera collina and Nephrama resupinatum. Specimens:

Occurrence Number: 008 Survey Site Name: ISABELLA CREEK County: Clearwater USGS quadrangle: SHEEP MTN. First Observed: 1995-06-21 Last Observed: 1995-06-21 Population Data: 1995: No data. Observation by Sarah Walker, Clearwater NF, and Juanita Lichthardt, Idaho CDC. Habitat Description: On branches in Thuja plicata/Athyrium filix-femina forest. Other lichens present include Lobaria pulmonaria. Specimens:

Occurrence Number: 009 Survey Site Name: AQUARIUS County: Clearwater USGS quadrangle: THOMPSON POINT First Observed: 1995-06-01 Last Observed: 1995-06-01 Population Data: 1995: No data. Collected by Sarah Walker, Clearwater NF. Habitat Description: NE aspect. Young Abies grandis stand with small Thuja plicata and Clintonia uniflora. On Thuja plicata branch with Lobaria pulmonaria. Other lichens present include Nephroma resupinatum, Hypogymnia physodes, Peltigera spp., and Alectora spp. Specimens: Sarah Walker 29 (herbarium unknown).

TEXOSPORIUM SANCTI-JACOBI

WOVENSPORE LICHEN
Occurrence Number: 001
Survey Site Name: SOUTH COLE ROAD/TENMILE CREEK SOUTH
County: Ada
USGS quadrangle: MORA
First Observed: 1993-05-18 Last Observed: 1993-05-18
Population Data: 1993: A scattered population of at least 50 clumps over 120 acres was assessed as having good population vigor by Anne DeBolt, Boise District BLM, on a cursory survey of the area on 5-18.
Habitat Description: These scattered populations were found on a gently sloping terrain with loamy-silt loam type soils.
Associated species include Artemisia tridentata wyomingensis, Poa secunda, Sitanion hystrix, and Stipa thurberiana. The overall site quality was rated as good.
Specimens: A. DeBolt 1679 (Boise BLM).

Occurrence Number: 002

Survey Site Name: CINDER CONE BUTTE

County: Ada

USGS quadrangle: CINDER CONE BUTTE

First Observed: 1993-05-11 Last Observed: 1995-01-17

Population Data: 1993: At least 20 clumps of lichen were observed. Unknown age class structure. Apothecia were present. Population vigor assessed as good. Observation on a cursory visit on 5-11 by Ann DeBolt, Boise District BLM. 1995: Ca 70-100 additional clumps with apothecia present were located to the SE, further up the NW slope of Cinder Cone Butte. Observation by Dana Quinney and J. Weaver, Idaho Army National Guard. Habitat Description: Occurs in a nice stand of Artemisia tridentata wyomingensis/Poa sandbergii/Stipa thurberiana habitat on

large volcanic cinders on the surface in some open areas. On clumps of dead Poa and mosses. Few exotic annuals present. Specimens: Ann DeBolt 1673 (Boise BLM).

Occurrence Number: 003

Survey Site Name: EIGHTMILE CREEK/TENMILE CREEK

County: Ada USGS quadrangle: CLOVERDALE

First Observed: 1991 Last Observed: 1993-05-18

Population Data: 1991: Hundreds of clumps. Population vigor assessed as high. Observation on a cursory visit by Ann DeBolt, Boise District BLM. 1993(Sec 18): At least 30 separate mature clumps were observed on a cursory visit by DeBolt. Population vigor assessed as good.

Habitat Description: Occurs in remnant Artemisia tridentata wyomingensis/native bunchgrass community on clumps of dead Poa. Associated with Sitanion hystrix, Stipa thurberiana, Agropyron spicatum, Poa sandbergii, and Lepidium papilliferum. On gently sloping terrain with northerly aspect. Specimens:

Occurrence Number: 004 Survey Site Name: EIGHTMILE CREEK County: Ada USGS quadrangle: BOISE SOUTH First Observed: 1993-07-27 Last Observed: 1993-07-27 Population Data: 1993: At least 15 clumps scattered over 40 acres. Population vigor assessed as moderate. Observation on a

cursory visit by Ann DeBolt, Boise District BLM. Habitat Description: Occurs on dead clumps of Poa secunda. Associates include Sitanion hystrix, Poa secunda, Agropyron spicatum, Stipa thurberiana, and Bromus tectorum. Site burned within the last 15 years so there are virtually no shrubs left. North aspect slope. Overall site quality rated as fair.

Specimens: Ann DeBolt 1755 (Boise BLM).

Occurrence Number: 005 Survey Site Name: KUNA BUTTE EAST County: Ada USGS quadrangle: KUNA First Observed: 1994-04-28 Last Observed: 1994-04-28 Population Data: 1994: 5 clumps. Population vigor assessed as fair. Observation by Ann DeBolt, Boise District BLM, on a cursory visit on 4-28. Habitat Description: Artemisia tridentata wyomingensis, Sitanion hystrix, Poa secunda site with good cryptogamic cover. Overall site quality rated as good. Specimens:

Occurrence Number: 006 Survey Site Name: ORCHARD SOUTHWEST County: Ada USGS quadrangle: ORCHARD First Observed: 1994-01-07 Last Observed: 1994-01-07 Population Data: 1994: Ca 1000 clumps with apothecia present. Observation by Dana Quinney and J. Weaver, Idaho Army National Guard. Habitat Description: Flat to gently rolling with fine silt soil. Weedy site with scattered sage. Associated species include Artemisia tridentata wyomingensis, Poa sandbergii, Ranunculus testiculatus, exotic Brassica spp., and Lepidium papilliferum. Also present in burned area next to the sage. Specimens: Occurrence Number: 007 Survey Site Name: HIGBY CAVE EAST - RED TIE AREA County: Ada USGS quadrangle: ORCHARD First Observed: 1993-12-03 Last Observed: 1993-12-03 Population Data: 1993: Several thousand clumps with apothecia present. Observation by Dana Quinney and J. Weaver, Idaho Army National Guard. Habitat Description: Slightly rolling terrain with fine silt soil. Growing on dead Poa clumps. Associated species include Artemisia tridentata wyomingensis, Poa sandbergii, Stipa thurberiana, and Lepidium papilliferum. Few exotic annuals are present. Specimens: Occurrence Number: 008 Survey Site Name: COLUMBIA VILLAGE - SOUTHEAST BOISE County: Ada USGS quadrangle: BOISE SOUTH Last Observed: 1994-06-24 First Observed: 1994-06-24 Population Data: 1994: Ca 30 clumps. Population vigor assessed as high. Area thoroughly surveyed by Ann DeBolt, Boise District BLM. Habitat Description: On clumps of humus in a patch of native bunchgrasses. Associates include Stipa thurberiana, Agropyron spicatum, Sitanion hystrix, and Poa secunda. Specimens: Occurrence Number: 009 Survey Site Name: PLEASANT VALLEY NORTH County: Ada USGS quadrangle: OWYHEE, IDAHO First Observed: 1987 Last Observed: 1987 Population Data: 1987: Ca 100+ clumps observed on a thorough survey by Ann DeBolt, Boise District BLM. Habitat Description: On humus clumps and soil. Artemisia tridentata wyomingensis-Stipa thurberiana-Agropyron spicatum community. Lepidium papilliferum is also present. Specimens: Roger Rosentreter 6815 (personal herbarium). Occurrence Number: 010 Survey Site Name: OWYHEE SOUTHWEST County: Ada USGS quadrangle: OWYHEE, IDAHO First Observed: 1983-04-28 Last Observed: 1988 Population Data: 1983: Hundreds of clumps along the E side of Pleasant Valley Road. Population vigor assessed as good to high. Observation by Roger Rosentreter, Idaho BLM. 1988: Hundreds of clumps (<1000) W of Pleasant Valley Road. Population vigor assessed as high. Observation on a cursory visit by Ann DeBolt, Boise District BLM.

Habitat Description: Flat area. Artemisia tridentata-Sitanion hystrix-Poa sandbergii and Chrysothamnus nauseosus-Poa secunda communities. Area intermixed with exotics Bromus tectorum and Sisymbrium altissimum. Specimens: Roger Rosentreter 3081 (Boise BLM, personal herbarium).

Occurrence Number: 011 Survey Site Name: SIMCO ROAD/I-84 County: Elmore USGS quadrangle: MAYFIELD SW First Observed: 1988 Last Observed: 1988 Population Data: 1988: Ca 50-100 scattered clumps. Population vigor assessed as high. Area thoroughly surveyed by Ann DeBolt, Boise District BLM, and Roger Rosentreter, Idaho BLM. Habitat Description: Flat topography. On soil and humus. Artemisia tridentata-Stipa thurberiana-Agropyron spicatum community in excellent condition. Lepidium papilliferum also present. Specimens: Roger Rosentreter 6819b (Boise BLM, personal herbarium). Bruce McCune 18894 (personal herbarium).

Occurrence Number: 012 Survey Site Name: MOUTH OF ISAACS CANYON County: Ada USGS quadrangle: BOISE SOUTH First Observed: 1995-10-11 Last Observed: 1995-10-11 Population Data: 1995: Ca 20-40 clumps observed on a cursory visit by Ann DeBolt, Lower Snake River District BLM. Population vigor assessed as high. Habitat Description: Portions of this general area are represented by a good to excellent bunchgrass (Agropyron spicatum, Stipa thurberiana, Sitanion hystrix)-Artemisia tridentata community. Some portions have no Artemisia tridentata left due to fire, but still have good bunchgrass representation. Weedier areas are intermittent (ie. dominated by Sisymbrium altissimum and Bromus tectorum). Specimens:

Occurrence Number: 013 Survey Site Name: N OF INITIAL POINT County: Ada USGS quadrangle: KUNA, INITIAL POINT First Observed: 1995-03 Last Observed: 1995-03 Population Data: 1995: 2 sites with a total of ca 30-50 clumps. Population vigor assessed as good. Observation on a cursory visit by Ann DeBolt, Boise District BLM. Habitat Description: Artemisia tridentata wyomingensis-Poa sandbergii-Sitanion hystrix community on the edge of a burn. Specimens:

THAMNOLIA VERMICULARIS

WORM LICHEN Occurrence Number: 001 Survey Site Name: ACROSS FROM HOSPITAL BAR County: Lemhi USGS quadrangle: RAMEY HILL First Observed: 1987-07-22 Last Observed: 1987-07-22 Population Data: 1987: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Moss-covered, N-facing small cliff band. Specimens: Roger Rosentreter 4525 (SRP).

XANTHOPARMELIA IDAHOENSIS

IDAHO RANGE LICHEN Occurrence Number: 001 Survey Site Name: LEMHI RIVER - SE OF MULKEY CREEK County: Lemhi USGS quadrangle: SAL MOUNTAIN First Observed: 1993-11-19 Last Observed: 1993-11-19 Population Data: 1993: Thousands of genets of unknown age class

Population Data: 1993: Thousands of genets of unknown age class structure were found at two sites. Population vigor assessed as very good in the larger of the two populations. Observation by Helen Ulmschneider, Salmon District BLM, who notes that more bare hills with potential habitat exist in the immediate area.

Habitat Description: Found on bare bentonite hills and flats bordering 2 small dry washes. Surrounding vegetation includes Artemisia tridentata and Sarcobatus vermiculatus. Other lichen include X. norchlorochroa on bare clay. Up to 80% of lichens were on south-facing hills above a small creek, but only 5-20% were on flats or north slopes. Overall lichen cover on all bare hills was roughly 10-20%, in patches. The overall site quality was rated as excellent.

Specimens: H. Ulmschneider 93-13 (Salmon BLM). H. Ulmschneider 93-12 (Salmon BLM).

Occurrence Number: 002 Survey Site Name: E OF SALMON AIRPORT - SE OF SALMON County: Lemhi USGS quadrangle: EAST OF SALMON First Observed: 1993-10-13 Last Observed: 1993-10-20

Population Data: 1993: 3 small populations of lichen were found. The easternmost population, visited on 10-14, had an estimated 2000 thalli clumps all of which were vegetative; population vigor assessed as good as about 5% of the X. Thalli present were X. idahoensis. The westernmost population, visited on 10-20, contained several hundred clumps; population vigor assessed as good, having ca 1-3% of lichens being X. idahoensis. The middle population, also visited on 10-20, contained ca 100 clumps; population vigor assessed as fair as <1% of the lichens present were X. idahoensis. Observations on cursory visits on 10-14 by Roger Rosentreter, Idaho State BLM, and Helen Ulmschneider, Salmon District BLM, and again on 10-20 by just Ulmschneider.

Habitat Description: Eastern population occurs on bentonite hills completely bare of vegetation in lacustrine ash deposits >40% clay. Slope <50% with generally southern aspects, but some also with northerly aspect. X. idahoensis occurs mixed in a large population of X. norchlorochroa, but it is easily distinguished from about 6 feet away. Surrounding vascular vegetation includes Atriplex spinosa, Tetradymia canescens. The western population also occurs on a south-facing, bare Bentonite slope with X. norchlorochroa and with Atriplex spinosa, Tetradymia spinescens, and Poa secunda. The middle population occurs on a rocky slope of small (1-3") stones with bentonite matrix. Also associated with X. norchlorochroa. Specimens: Roger Rosentreter 8238 (SRP, US, Salmon BLM). Helen Ulmschneider 93-12 (Salmon BLM). Helen Ulmschneider 93-13 (Salmon BLM).

Occurrence Number: 003 Survey Site Name: HOT SPRING CREEK County: Lemhi USGS quadrangle: SAL MOUNTAIN First Observed: 1986-01-04 Last Observed: 1986-01-04 Population Data: 1986: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: On calcareous lacustrine ash soil. Sites nearly barren of vascular vegetation and surrounded by Atriplex confertifolia. Specimens: Roger Rosentreter 3828 (Personal herbarium, isotype; US, holotype).

Occurrence Number: 004 Survey Site Name: KADLETZ CREEK County: Lemhi USGS quadrangle: BAKER First Observed: 1987-07-07 Last Observed: 1987-07-07 Population Data: 1987: No data. Collected by Roger Rosentreter, Idaho State BLM. Habitat Description: Barren benthene clay soil, Artemisia tridentata wyomingensis and Atriplex habitat. Specimens: Roger Rosentreter 4520 (SRP) - determined by M. Hale 1987.

Occurrence Number: 005 Survey Site Name: N OF PRICE CREEK County: Lemhi USGS quadrangle: BAKER First Observed: 1995-05-19 Last Observed: 1995-05-19 Population Data: 1995: 3 populations. The top of hill 4773T is almost a solid carpet of X. idahoensis. Population vigor assessed as excellent. Observation on a cursory visit by Helen Ulmschneider, Salmon District BLM, noting that there are extensive clay hills in the area that were not visited. Habitat Description: Growing on clay. Bare clay hills with Monolepsis nuttalliana, Tetradymia sp., and Artemisia sp. Other range lichens present include Xanthoparmelia norchlorochroa and X. wyomingensis in the shrub area.

Specimens: Helen Ulmschneider 95-01, 95-02, 95-03 (Salmon BLM).

CONCLUSIONS AND RECOMMENDATIONS

1. **GLOBALLY RARE SPECIES** - While most of our state's rare lichens and bryophytes are globally common, there are several species that are globally rare (*i.e.*, have a global conservation rank of G1, G2, or G3 in Tables 1 and 3). These species should be of particular conservation concern to land managers and biologists in the state. The globally rare species occurring in Idaho are:

Bryophytes

Bryum calobryoides G3 Orthotrichum holzingeri G1 or G2? Sphaerocarpos hians G1

Lichens

Catapyrenium congestum G2 Cetraria subalpina G3 Cladonia andreggii G1 Cladonia luteoalba G2 Collema curtisporum G1 Dermatocarpon lorenzianum G1 Texosporium sancti-jacobi G2 Xanthoparmelia idahoensis G1

2. **INCREASE AWARENESS FOR RARE CRYPTOGAMS** - Rare bryophytes and lichens should be incorporated into the overall rare species and ecosystem conservation activities of land management agencies and organizations, similar to vascular plants and vertebrate animals. The first step is to increase the ability of biologists to identify and inventory for rare species through special training and workshops. While a few training sessions have taken place in the last couple of years, more are needed.

The results of rare cryptogam inventories should be submitted to the CDC so that populations can be entered into our rare species occurrence data base, which is the centralized data base for Idaho. Reporting forms and instructions can be downloaded from the CDC's home page on the World Wide Web (address: www.state.id.us/fishgame/cdchome.htm)

3. **RARE FUNGI -** The diversity of fungi in Idaho is particularly notable, from the rich diversity in the central Idaho mountains to the endemism on the Owyhee Plateau. Although technically not plants, but usually lumped in with the field of botany, this group of organisms are important members of our native ecosystems. Only recently has the conservation of fungi species begun to be addressed in the Pacific Northwest, in general, and Idaho in particular. Information on rare fungi compiled as part of the Interior Columbia River Basin assessment needs to be incorporated into the CDC's data base and into species and ecosystem conservation activities of land management agencies.

4. **NONVASCULAR WORKING GROUP** - We recommend that a "working group" be established to evaluate the conservation of nonvascular plants in the state. This is usually done at the annual Idaho Rare Plant Conference, but for several reasons (listed below) we believe that a separate group could more effectively evaluate the status of these species: 1) considerable time needs to be spent on the suggested additions to the bryophyte list and the nearly *de novo* evaluation of rare fungi in the state; 2) the hectic schedule of the conference rarely (never?) allows for such an in depth evaluation; and 3) relatively few people have adequate knowledge of nonvcascular groups. This group clearly needs to address several short-term issues and may not need to be permanent.

BRYOPHYTE AND LICHEN REFERENCES AT THE CDC

Alaback, P. B. 1982. Dynamics of understory biomass in Sitka spruce-western hemlock forests of southeast Alaska. Ecology 63(6): 1932-1948.

RUBUS SPECTABILIS MONTANE FOREST OLD GROWTH FOREST THELYPTERIS PHEGOPTERIS PHEGOPTERIS CONNECTILIS HOOKERIA LUCENS BLECHNUM SPICANT HYPOGYMNIA ENTEROMORPHA

Alstrup, V., and U. Sochting. 1989. Checkliste og status over Danmarks laver. Nordisk Lichenologisk Forening. Kobenhavn, Denmark. 44 pp.

LICHEN CHECKLIST COLLEMA FURFURACEUM CLADONIA UNCIALIS LOBARIA SCROBICULATA PHYSCIA SEMIPINNATA

Anderegg, D. E., G. J. Schroeder, and N. E. Schroeder. 1973. Further additions to the lichen flora of Idaho. The Bryologist 76(1): 207-208.

LICHEN DISTRIBUTION IDAHO

Andrus, R. E., and E. F. Layser. 1976. Sphagnum in the northern Rocky Mountains of the United States. The Bryologist 79: 508-511.

SPHAGNUM MENDOCINUM DISTRIBUTION CHECKLIST

Andrus, R. E. 1989. Letter dated August 25, 1989, requesting information on rare and endangered bryophytes in order to develop a North American database.

NONVASCULAR CONSERVATION

Andrus, R. E., E. F. Karlin, and S. S. Talbot. 1992. Rare and endangered Sphagnum species in North America. Biological Conservation 59: 247-254.

SPHAGNUM MENDOCINUM NONVASCULAR CONSERVATION DISTRIBUTION

Berg, A., B. Ehnstrom, L. Gustafsson, T. Hallingback, M. Jonsell, and J. Weslien. 1995. Threat levels and threats to red-listed species in Swedish forests. Conservation Biology 9(6): 1629-1633.

LICHEN BRYOPHYTE NONVASCULAR CONSERVATION

Bird, C. D., and A. H. Marsh. Phytogoegraphy and ecology of the lichen family Parmeliaceae in southwestern Alberta. Canadian Journal of Botany 51: 261-288.

HYPOGYMNIA ENTEROMORPHA PHYTOGEOGRAPHY DISTRIBUTION

Bohlin, A., L. Gustafsson, and T. Hallingback. 1977. Skirmossan, Hookeria lucens, i Sverige. Svensk Bot. Tidskr 71: 273-284.

HOOKERIA LUCENS DISTRIBUTION COMMUNITY ECOLOGY SPECIES BIOLOGY MANAGEMENT TECHNIQUES

Breuss O., and B. McCune. 1994. Addition to the pyrenolichen flora of North America. The Bryologist 97(4) pp. 365-370.

CATAPYRENIUM CONGESTUM DERMATOCARPON LORENZIANUM LICHEN DISTRIBUTION TAXONOMY

Broad, K. 1989. Lichens in southern woodlands. HMSO Books, London, England. 48 p.

LOBARIA SCROBICULATA LICHEN

Chadde, S., and S. Shelly. 1994. Significant peatlands of western Montana: site descriptions and major features. Unpublished report prepared for USDA Forest Service, Northern Region. 41 pp. plus appendices.

PEATLAND INVENTORY CLASSIFICATION BRYOPHYTE CHECKLIST SPHAGNUM MENDOCINUM

Christy, J. A., J. H. Lyford, and D. H. Wagner. 1982. Checklist of Oregon mosses. The Bryologist 85(1): 22-36.

SPHAGNUM MENDOCINUM TAYLORIA SERRATA ULOTA MEGALOSPORA HOOKERIA LUCENS HELODIUM BLANDOWII BUXBAUMIA APHYLLA BUXBAUMIA PIPERI CHECKLIST

Christy, J. A., and J. S. Harpel. 1995. Bryophytes of the Columbia River Basin, south of the Canadian border.

Unpublished report prepared for the Eastside Ecosystem Management Project. 48 pp. plus appendices.

BRYOPHYTE NONVASCULAR CONSERVATION BIOGEOGRAPHY FUNCTION BRYUM CALOBRYOIDES BUXBAUMIA APHYLLA BUXBAUMIA PIPERI HELODIUM BLANDOWII HOOKERIA LUCENS MEESIA LONGISETA SPHAGNUM MENDOCINUM TAYLORIA SERRATA SPHAEROCARPOS HIANS

Clark, L. and T. C. Frye. 1928. Liverworts of the Northwest. Publications of the Puget Sound Biological Station 6: 1-94. (p. 28-30).

SPHAEROCARPOS HIANS TAXONOMY DISTRIBUTION BRYOPHYTE

Cooke, W. B. 1955. Fungi, lichens and mosses in relation to vascular plant communities in eastern Washington and adjacent Idaho. Ecological Monographs 25(2): 119-180.

MONTANE FOREST WOODLAND GRASSLAND FUNGI LICHEN BRYOPHYTE CLASSIFICATION

Corner, R. W. M. 1967. Further observations on the distribution of Buxbaumia aphylla Hedw. on colliery debris. Transactions of the British Bryological Society 5: 828-829.

BUXBAUMIA APHYLLA COMMUNITY ECOLOGY SPECIES BIOLOGY

Coxson, D. S., G. P. Harris, and K. A. Kershaw. 1982. Physiological-environmental interactions in lichens. XV. Contrasting gas exchange patterns between a lichenized and non-lichenized terrestrial Nostoc cyanophyte. New Phytologist 92: 561-572.

COLLEMA FURFURACEUM SPECIES BIOLOGY

Cros i Matas, R. M. 1983. Tayloria serrata (Hedw.) B.S.G. a la Cerdyana. Collectanea Botanica "a Barcionesi Botanico Instituto Edita" 14: 261-263.

DISTRIBUTION TAYLORIA SERRATA

DeBolt, A., and B. McCune. 1993. Lichens of Glacier National Park, Montana. The Bryologist 96(2): 192-204.

LOBARIA HALLII HYPOGYMNIA ENTEROMORPHA CHECKLIST COLLEMA CURTISPORUM COLLEMA FURFURACEUM

DeBolt, A. Habitat conservation assessment for slick-spot peppergrass (Lepidium papilliferum) and woven-spore lichen (Texosporium sancti-jacobi). Unpublished draft report. 13 pp. plus appendices.

STATUS SURVEY LEPIDIUM PAPILLIFERUM TEXOSPORIUM SANCTI-JACOBI

Dewey, R. M. 1983. Bryophytes of the Tenmile Creek drainage, Idaho. The Bryologist 86(2): 114-116.

BRYOPHYTE CHECKLIST

Duke, T. 1989. Save the Cairngorms campaign. British Lichen Society Bulletin 65: 34-35.

NONVASCULAR CONSERVATION

Elliott, J. C. 1989. Rare mosses of Idaho. Unpublished report. 3 pp.

MULTI-SPECIES NONVASCULAR CONSERVATION BRYUM CALOBRYOIDES BUXBAUMIA PIPERI BUXBAUMIA APHYLLA HELODIUM BLANDOWII HOOKERIA LUCENS MEESIA LONGISETA SPHAGNUM MENDOCINUM TAYLORIA SERRATA ULOTA CURVIFOLIA ULOTA MEGALOSPORA

Elliott, J. C., and G. L. Moore. 1989. Additions to the moss flora of Montana. The Bryologist 92(2): 194-197.

SPHAGNUM MENDOCINUM DISTRIBUTION

Elliott, J. C. 1989. Rare mosses of Idaho. Unpublished report. 3 pp.

MULTI-SPECIES NONVASCULAR CONSERVATION BRYUM CALOBRYOIDES BUXBAUMIA PIPERI BUXBAUMIA APHYLLA HELODIUM BLANDOWII HOOKERIA LUCENS MEESIA LONGISETA SPHAGNUM MENDOCINUM TAYLORIA SERRATA ULOTA CURVIFOLIA ULOTA MEGALOSPORA Eversman, S., and L. l. Sigal. 1984. Ultrastructural effects of peroxyacetyl nitrate (PAN) on two lichen species. The Bryologist 87(2): 112-118.

HYPOGYMNIA ENTEROMORPHA SPECIES BIOLOGY

Eversman, S. 1990. Lichens of Yellowstone National Park. The Bryologist 93(2): 197-205.

YELLOWSTONE FNPNP LICHEN CHECKLIST

Eversman, S. 1994. Lichens of the Yellowstone eocsystem. Unpublished report prepared for the East Side Project, Columbia River Basin Ecosystem Management Plan. On file at Idaho Department of Fish and Game, Conservation Data Center, Boise. 93 pp.

YELLOWSTONE FNPNP GYE LICHEN DISTRIBUTION COMMUNITY ECOLOGY THAMNOLIA VERMICULARIS HYPOGYMNIA ENTEROMORPHA

Foster, D. R., H. E. Wright, Jr., M. Thelaus, and G. A. King. 1988. Bog development and lanfdorm dynamics in central Sweden and south-eastern Labrador, Canada. Journal of Ecology 76: 1164-1185.

FUNCTION PEATLAND VEGETATION HISTORY BRYOPHYTE HYDROLOGY VACCINIUM OXYCOCCOS SCIRPUS CAESPITSOUS

Geiser, L. H., C. C. Derr, and K. L. Dillman. 1994. Air quality monitoring on the Tongass National Forest: Methods and baselines using lichens. USDA Forest Service R10-TB-46. Tongass National Forest, Stikine Area, Petersburg, AK. 84 pp. plus appendices.

AIR QUALITY THAMOLIA VERMICULARIS PILOPHORUS ACICULARIS CLADONIA TRANSCENDENS CLADONIA UNCIALIS COLLEMA FURFURACEUM PSEUDOCYPHELLARIA ANTHRASPIS HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA LINITA LOBARIA SCROBICULATA

Geiser, L. 1994. Why sample lichens? Unpublished report prepared for the Columbia River Basin Project, Science Integration Team. Not paged.

AIR QUALITY LICHEN

Geiser, L. H., C. C. Derr, and K. L. Dillman. 1994. Air quality monitoring on the Tongass National Forest: Methods and baselines using lichens. USDA Forest Service R10-TB-46. Tongass National Forest, Stikine Area, Petersburg, AK. 84 pp. plus appendices.

AIR QUALITY THAMOLIA VERMICULARIS PILOPHORUS ACICULARIS CLADONIA TRANSCENDENS CLADONIA UNCIALIS COLLEMA FURFURACEUM PSEUDOCYPHELLARIA ANTHRASPIS HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA LINITA LOBARIA SCROBICULATA

Gignac, L. D., D. H. Vitt, S. C. Zoltai, and S. E. Bayley. 1991. Bryophyte response surfaces along climatic, chemical, and physical gradients in peatlands of western Canada. Nova Hedwigia 53: 27-71.

PEATLAND BRYOPHYTE

Goffinet, B. 1994. Floristic notes on the lichenicolous fungi flora of Alberta, Canada. Mycotaxon LI: 1-4.

DISTRIBUTION LOBARIA HALLII

Gough, L. P., L. L. Jackson, and J. A. Sacklin. 1988. Determining baseline element composition of lichens. II. Hypogymnia enteromorpha and Usnea spp. at Redwood National Park, California. Water, Air, and Soil Pollution 38: 169-180.

HYPOGYMNIA ENTEROMORPHA SPECIES BIOLOGY

Goward, T. 1990. Rare and endangered lichens in B.C. ABB BioLine (Official publication of the Association of Professional Biologists of British Columbia) 9(2): 13-15.

NONVASCULAR CONSERVATION

Goward, T. 1991. Epiphytic lichens: Going down with the trees. Pages 153-158 in S. Rautio, editor, Community action for endangered species: A public symposium on B.C.'s threatened & endangered species and their habitat. The Federation of British Columbia Naturalists and Northwest Wildlife Preservation Society. Vancouver, B.C.

OLD GROWTH FOREST MONTANE FOREST NONVASCULAR CONSERVATION HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA LINITA PSEUDOCYPHELLARIA ANTHRASPIS

Goward, T., P. Diederich, and R. Rosentreter. 1994. Notes on the lichens and allied fungi of British Columbia. II. The Bryologist 97(1): 56-62.

CLADONIA LUTEOALBA DISTRIBUTION

Hale, M. E. 1989. New species in the lichen genus Xanthoparmelia (Ascomycotina: Parmeliaceae). Mycotaxon 34(4): 541-564.

XANTHOPARMELIA IDAHOENSIS

Hale, M. E. 1990. A synopsis of the lichen genus Xanthoparmelia (Vianio) Hale (Ascomycotina, Parmeliaceae). Smithsonian Contributions to Botany, number 74. 250 pp.

XANTHOPARMELIA IDAHOENSIS TAXONOMY

Hallingback, T. 1989. Occurrence and ecology of the lichen Lobaria scrobiculata in southern Sweden. Lichenologist 21(4): 331-341.

LOBARIA SCROBICULATA NONVASCULAR CONSERVATION DISTRIBUTION

Hammer, S., and T. Ahti. 1990. New and interesting species of Cladonia from California. Mycotaxon 37: 335-348.

LICHEN DISTRIBUTION CLADONIA VERRUCULOSA

Hammer, S. 1995. A synopsis of the genus Cladonia in the northwestern United States. The Bryologist 98(1): 1-28.

TAXONOMY BIOGEOGRAPHY CLADONIA IMBRICARICA CLADONIA UNCIALIS CLADONIA LUTEOALBA CLADONIA TRANSCENDENS CLADONIA ANDEREGGII

Hammer, S. 1995. The biogeography and ecology of species in the lichen genus Cladonia in the Columbia River Basin. Unpublished report prepared for the Eastside Ecosystem Management Project. 65 pp.

CLADONIA UNCIALIS CLADONIA TRANSCENDENS CLADONIA ANDEREGGII CLADONIA LUTEOALBA CLADONIA IMBRICARICA COMMUNITY ECOLOGY BIOGEOGRAPHY DISTRIBUTION

Hammon, D., and L. C. Pearson. 1967. Lichens of eastern Idaho and adjacent Wyoming, Utah and Montana in the Ricks College Herbarium. Journal of the Idaho Academy of Science 12(2): 55-57.

CHECKLIST IDAHO HYPOGYMNIA ENTEROMORPHA Hancock, J. A., and G. R. Brassard. 1974. Phenology, sporophyte production, and life history of Buxbaumia aphylla in Newfoundland, Canada. The Bryologist 77(4): 501-513.

BUXBAUMIA APHYLLA SPECIES BIOLOGY

Hancock, J. A., and G. R. Brassard. 1974. Element content of moss sporophytes: Buxbaumia aphylla. Canadian Journal of Botany 52: 1861-1865.

BUXBAUMIA APHYLLA SPECIES BIOLOGY

Hayward, G. D., and P. H. Hayward. 1988. Nest box use: Idaho Panhandle National Forest, 1988. University of Idaho, Department of Fish and Wildlife Resources, College of Forestry, Wildlife, and Range Sciences, Moscow. 4 pp.

IDAHO PANHANDLE NATIONAL FORESTS BOREAL OWL NORTHERN SAW-WHET OWL RAPTORS REPRODUCTION CAVITY NESTERS AEGOLIUS ACADICUS AMERICAN KESTREL FALCO SPARVERIUS NORTHERN FLYING SQUIRREL GLAUCOMYS SABRINUS AEGOLIUS FUNEREUS LICHEN

Hayward, G. D., and P. H. Hayward. 1988. Nest box use: Payette National Forest, 1988. University of Idaho, Department Fish and Wildlife Resources, College of Forestry, Wildlife, and Range Sciences, Moscow. 4 pp.

PAYETTE NATIONAL FOREST RAPTORS BOREAL OWL NORTHERN SAW-WHET OWL AEGOLIUS ACADICUS CAVITY NESTERS NORTHERN FLYING SQUIRREL GLAUCOMYS SABRINUS MARTEN MARTES AMERICANA AEGOLIUS FUNEREUS LICHEN REPRODUCTION BIOLOGY

Hayward, G. D., and R. Rosentreter. 1994. Lichens as nesting material for northern flying squirrels in the northern Rocky Mountains. Journal of Mammalogy 75(3): 663-673.

SPECIES BIOLOGY GLAUCOMYS SABRINUS NORTHERN FLYING SQUIRREL SMALL MAMMALS LICHEN

PAYETTE NATIONAL FOREST

Hoffman, G. R. 1974. The influence of a paper pulp mill on the ecological distribution of epiphytic cryptogams in the vicinity of Lewiston, Idaho, and Clarkston, Washington. Environmental Pollution 7: 283-301.

IDAHO AIR QUALITY LICHEN

Holthausen, R. and C. Meslow, technical coordinators. 1993. Terrestrial forest ecosystem assessment. Chapter 4 in J. W. Ward et al., Forest ecosystem management: An ecological, economic, and social assessment. USDA Forest Service, USDI Bureau of Land Management, National Park Service, and Fish and Wildlife Service, USDC National Marine Fisheries Service, and the Environmental Protection Agency. Washington DC.

NONVASCULAR CONSERVATION MONTANE FOREST BUXBAUMIA PIPERI HOOKERIA LUCENS RAMALINA THRAUSTA PSEUDOCYPHELLARIA ANTHRASPIS LOBARIA HALLII LOBARIA SCROBICULATA LOBARIA LINITA

Jamieson, D. W., and L. G. Holmberg. 1969. Buxbaumia piperi new to California. The Bryologist 72: 72-73.

BUXBAUMIA PIPERI DISTRIBUTION

Jamieson, D. W. 1989. The bryoflora of the southern Rocky Mountains of Colorado. American Journal of Botany 76: 9. Abstract.

BUXBAUMIA PIPERI DISTRIBUTION

Jones, E. W. [1968?]. The ecology of Buxbaumia aphylla Hedw. Transactions of the British Bryological Society 6: 139.

BUXBAUMIA APHYLLA COMMUNITY ECOLOGY SPECIES BIOLOGY

Jordan, W. P. 1973. The genus Lobaria in North America north of Mexico. The Bryologist 76(2): 225-251.

LOBARIA HALLII LOBARIA LINITA LOBARIA SCROBICULATA TAXONOMY

Kershaw, K. A. 1974. Studies on lichen-dominated systems. X. The sedge meadows of the coastal raised beaches. Canadian Journal of Botany 52: 1947-1972.

LICHEN COMMUNITY ECOLOGY SALIX CANDIDA ANDROMEDA POLIFOLIA

CAREX CHORDORRHIZA

Kershaw, K. A., and J. D. MacFarlane. 1982. Physiological-environmental interactions in lichens. XIII. Seasonal constancy of nitrogenase activity, net photosysthesis and respiration, in Collema furfuraceum (Am.) Dr. New Phyotologist 90: 723-734.

COLLEMA FURFURACEUM SPECIES BIOLOGY

Koponen, T. 1992. Endangered bryophytes on a global scale. Biological Conservation 59: 255-258.

NONVASCULAR CONSERVATION

Kristinsson, H. 1981. Additions to the lichen flora of Iceland II. Acta Botanica Islandica 6: 23-28.

COLLEMA FURFURACEUM DISTRIBUTION

Krog, H., and P. W. James. 1977. The genus Ramalina in Fennoscandia and the British Isles. Norwegian Journal of Botany 24: 15-43.

RAMALINA THRAUSTA LICHEN TAXONOMY

Kuhry, P., B. J. Nicholson, L. D. Gignac, D. H. Vitt, and S. E. Bayley. 1993. Development of Sphagnum-dominated peatlands in boreal continental Canada. Canadian Journal of Botany 71: 10-22.

PEATLAND BRYOPHYTE VEGETATION HISTORY VACCINIUM OXYCOCCOS

Leonard, B. F., and R. Rosentreter. 1994. Dating a 20th-Century fault, Elk Summit talus apron, Big Creek area, Valley County, Idaho. U.S. Geological Survey Bulletin 2101. U.S. Government Printing Office, Washington D.C. 13 pp.

LICHEN GEOLOGY IDAHO

Lesher, R. D., R. Rosentreter, and J. Christy. 1994. The role of fungi, lichens, and bryophytes in the development of management alternatives for federal lands in the Pacific Northwest. Northwest Science 68(2): 97. Abstract.

MONTANE FOREST LICHEN BRYOPHYTE

Lesica, P., B. McCune, S. V. Cooper, and W. S. Hong. 1991. Differences in lichen and bryophyte communities between old-growth and managed second-growth forests in the Swan Valley, Montana. Canadian Journal of Botany 69: 1745-1755.

OLD GROWTH FOREST MONTANE FOREST COMMUNITY ECOLOGY MANAGEMENT TECHNIQUES BRYOPHYTE LICHEN HYPOGYMNIA ENTEROMORPHA

Lichthardt, J. 1995. Summary of lichen and moss EORs from collections at the University of Idaho and Washington State University herbaria.

MULTI-SPECIES BUXBAUMIA PIPERI HELODIUM BLANDOWII HOOKERIA LUCENS SPHAGNUM MENDOCINUM CETRARIA SUBALPINA HYPOGYMNIA APINNATA HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA SCROBICULATA PSEUDOCYPHELLARIA ANTHRASPIS

Lin, C. K., J. Y. Hsiao, and M. J. Lai. 1989. Studies on the lichen genus Lobaria in Taiwan. Proceedings of the National Science Council, Republic of China, Part B Life Sciences 13(1): 47-55.

LOBARIA SCROBICULATA TAXONOMY DISTRIBUTION

Maass, W. S. G. 1967. Studies on the taxonomy and distribution of Sphagnum IV: Sphagnum majus, Sphagnum annulatum, Sphagnum mendocinum, and Sphagnum obtusum in North America. Nova Hedwigia 14: 187-214.

SPHAGNUM MENDOCINUM TAXONOMY DISTRIBUTION

Maass, W. S. G. 1980. Erioderma pedicellatum in North America: A case study of a rare and endangered lichen. Proceedings of the Nova Scotia Institute of Science 30: 69-87.

NONVASCULAR CONSERVATION

MacFarlane, J. D., and K. A. Kershaw. 1982. Physiological-environmental interactions in lichens. XIV. The environmental control of glucose movement from alga to fungus in Peltigera polydactyla, P. rufescens and Collema furfuraceum. New Phytologist 91: 93-101.

COLLEMA FURFURACEUM SPECIES BIOLOGY

McCleary, J. A., and V. V. Green. 1971. A checklist of Idaho mosses. The Bryologist 74: 175-180.

CHECKLIST DISTRIBUTION TAYLORIA SERRATA MEESIA LONGISETA HOKERIA LUCENS BUXBAUMIA APHYLLA BUXBAUMIA PIPERI McCoy, P. R. 1994. Lichen (not moss) forest-health indicator. Capital Press, August 19, 1994, edition. Boise, ID.

LICHEN

McCoy, P. R. 1994. Botanist shows Polish scientists some tricks. Capital Press, August 19, 1994, edition. Boise, ID.

LICHEN

McCune, B. 1977. Vegetation development on a low elevation talus slope in western Montana. Northwest Science 51(3): 198-207.

LICHEN BRYOPHYTE CLASSIFICATION SUCCESSION MONTANE FOREST COMPOSITION STRUCTURE

McCune, B. 1982. Lichens of the Swan Valley, Montana. The Bryologist 85(1): 13-21.

COLLEMA CURTISPORUM COLLEMA FURFURACEUM LOBARIA HALLII PHYSCIA SEMIPINNATA CHECKLIST LICHEN

McCune, B. 1984. Lichens with oceanic affinities in the Bitterroot Mountains of Montana and Idaho. The Bryologist 87(1): 50-56.

LICHEN AQUARIUS FFSRN CLADONIA TRANSCENDENS HYPOGYMNIA ENTEROMORPHA PSEUDOCYPHELLARIA ANTHRASPIS SPHAEROPHORUS TUCKERMANNII LOBARIA LINITA BIOGEOGRAPHY DISTRIBUTION

McCune, B., and R. Rosentreter. 1988. Proposed nonvascular sensitive plant species for the Northern Region of the USFS. Unpublished report on file at Conservation Data Center, Idaho Department of Fish and Game, Boise. 2 pp.

LOBARIA LINITA COLLEMA FURFURACEUM PHYSCIA SEMIPINNATA LOBARIA HALLII COLLEMA CURTISPORUM NONVASCULAR CONSERVATION DISTRIBUTION McCune, B. 1990. Rapid estimation of abundance of epiphytes on branches. The Bryologist 93(1): 39-43.

SAMPLING LICHEN MONTANE FOREST

McCune, B. 1991. Field key to the lichens of the Northwest forests west of the Cascade Crest. Unpublished report. 19 pp.

FIELD GUIDE LICHEN PSEUDOCYPHELLARIA ANTHRASPIS COLLEMA FURFURACEUM RAMALINA THRAUSTA

McCune, B., and A. DeBolt. 1991. Lichens from proposed Aquarius Research Natural Area. Unpublished list on file at the Conservation Data Center, Idaho Department of Fish and Game, Boise. 2 pp.

AQUARIUS FFSRN CLEARWATER NATIONAL FOREST SPHAEROPHORUS TUCKERMANNII RAMALINA THRAUSTA PSEUDOCYPHELLARIA ANTHRASPIS HYPOGYMNIA ENTEROMORPHA CLADONIA TRANSCENDENS

McCune, B. 1992. Lichens of southwest Idaho. Class 1 survey of lichen flora of southwest Idaho. Final Report. Cooperative Challange Cost Share Project, Bureau of Land management, Boise District Office, and McCune and Muir Ecological Services. 52 pp.

MUD FLAT OOLITE FBLAC CHECKLIST TEXOSPORIUM SANCTI-JACOBI LICHEN

McCune, B. 1992. Status of a globally ranked (G2) rare lichen species, Texosporium sancti-jacobi. Summary report of the status of the species and habitat management recommendations. Final Report. Cooperative Challange Cost Share Project, Bureau of Land Management, Boise District Office, and McCune and Muir Ecological Services. 38 pp.

TEXOSPORIUM SANCTI-JACOBI STATUS SURVEY

McCune, B., and R. Rosentreter. 1993. Improved spot tests for Bryoria. Evansia 10(2): 58-60.

LICHEN SPECIES BIOLOGY

McCune, B. 1994. Lichen database listing for Columbia Basin. Unpublished report prepared for the Eastside Ecosystem Management Project. 209 pp.

PHYSCIA SEMIPINNATA PILOPHORUS ACICULARIS TEXOSPORIUM SANCTI-JACOBI HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA SCROBICULATA LICHEN CHECKLIST ASPICILIA FRUTICULOSA CLADINA RANGIFERINA CLADONIA LUTEOALBA CLADONIA TRANSCENDENS CLADONIA UNCIALIS COLLEMA CURTISPORUM COLLEMA FURFURACEUM HYPOGYMNIA APINNATA

McCune, B. 1994. Lichen species groups in the Columbia Basin - ecosystem functions and indicator values. Unpublished report prepared for the Eastside Ecosystem Management Project. 28 pp.

LICHEN FUNCTION CHECKLIST ASPICILIA FRUTICULOSA CLADINA RANGIFERINA CLADONIA LUTEOALBA CLADONIA TRANSCENDENS CLADONIA UNCIALIS COLLEMA CURTISPORUM COLLEMA FURFURACEUM HYPOGYMNIA ENTEROMORPHA LOBARIA SCROBICULATA LOBARIA HALLII LOBARIA LINITA PHYSCIA SEMIPINNATA PILOPHORUS ACICULARIS **TEXOSPORIUM SANCTI-JACOBI** CATAPYRENIUM CONGESTUM

McCune, B., and R. Rosentreter. 1995. Distribution and ecology of Thelomma ocellatum in western North America. Evansia 12(3): 103-106.

THELOMMA OCELLATUM DISTRIBUTION LICHEN

McCune, B., and T. Goward. 1995. Macrolichens of the Northern Rocky Mountains. Mad River Press, Inc. Eureka, California. 208 pp.

CLADONIA UNCIALIS CLADONIA LUTEOALBA CLADONIA TRANSCENDENS CLADINA RANGIFERINA ASPICILIA FRUTICULOSA LICHEN FLORA CETARIA SUBALPINA COLLEMA CURTISPORUM COLLEMA FURFURACEUM HYPOGYMNIA APINNATA HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA LINITA LOBARIA SCROBICULATA PHYSCIA SEMIPINNATA PILOPHORUS ACICULARIS PSEUDOCYPHELLARIA ANTHRASPIS THAMNOLIA VERMICULARIS

McCune, B., R. Rosentreter, and A. Debolt. 1995. Lichens from Bottle Lake RNA. 1 p.

LICHEN BOTTLE LAKE FFSRN

McCune, B., and R. Rosentreter. 1995. Field key to soil lichens of central and eastern Oregon. Unpublished report. 9 pp.

FIELD GUIDE ASPICILIA FRUTICULOSA TEXOSPORIUM SANCTI-JACOBI

McCune, B. [n.d.]. Pacific Northwest epiphytic lichens: Species codes. Unpublished list on file at the Conservation Data Center, Idaho Department of Fish and Game, Boise. 1 pp.

LOBARIA SCROBICULATA LOBARIA LINITA LOBARIA HALLII HYPOGYMNIA ENTEROMORPHA CLADONIA TRANSCENDENS CLADONIA FURFURACEUM PHYSCIA SEMIPINNATA PSEUDOCYPHELLARIA ANTHRASPIS RAMALINA THRAUSTA

Miller, N. G. 1971. Three additions to the moss flora of New York State. The Bryologist 74: 377-378.

TAYLORIA SERRATA DISTRIBUTION

Moseley, R. K. 1995. The ecology of geothermal springs in south-central Idaho. Cooperative study, Sawtooth National Forest and Conservation Data Center, Idaho Department of Fish and Game. 47 pp. plus appendices.

EPIPACTIS GIGANTEA EPILOBIUM PALUSTRE PRIMULA INCANA WETLAND NONVASCULAR CONSERVATION YELLOWSTONE FNPNP HOT SPRINGS AQUATIC THERMAL SPRINGS AQUATIC COMMUNITY MANAGEMENT TECHNIQUES Multiple authors. 1993. The Bryophytes of peat-forming ecosystems. Bryological and lichenological section symposium, Supplement to the American Journal of Botany Abstracts 80(6): 1-3.

PEATLAND BRYOPHYTE VEGETATION HISTORY CLASSIFICATION

Nash, T. H., and L. L. Sigal. 1979. Gross photosynthetic response of lichens to short-term ozone fumigations. The Bryologist 82(2): 280-285.

HYPOGYMNIA ENTEROMORPHA AIR QUALITY SPECIES BIOLOGY

Nilsson, S. G., U. Arup, R. Baranowski, and S. Ekman. 1995. Tree-dependent lichens and beetles as indicators in conservation forests. Conservation Biology 9(5): 1208-1215.

LICHEN NONVASCULAR CONSERVATION

Noble, M. G., D. B. Lawrence, and G. P. Streveler. 1984. Sphagnum invasion beneath an evergreen forest canopy in southeastern Alaska. The Bryologist 87(2): 119-127.

COMMUNITY ECOLOGY SPHAGNUM MENDOCINUM

Owen, W. 1994. Sensitive Plant Species of the Boise National Forest. Boise National Forest, ID. 5 pp.

FIELD GUIDE MULTI-SPECIES ALLIUM TOLMIEI PERSIMILE BRYUM CALOBRYOIDES DOUGLASIA IDAHOENSIS HAPLOPAPPUS INSECTICRURIS PHACELIA MINUTISSIMA ALLIUM VALIDUM CAREX BUXBAUMII EPIPACTIS GIGANTEA LEWISIA KELLOGGII RHYNCHOSPORA ALBA SCHEUCHZERIA PALUSTRIS

Pickett, F. L., and G. N. Jones. 1928. Check-list of the mosses known to occur in southeastern Washington and adjacent Idaho. The Bryologist 31: 54-56.

BRYOPHYTE CHECKLIST

Pike, L. H., and M. E. Hale, Jr. 1982. Three new species of Hypogymnia from western North America (Lichenes Hypogymniaceae). Mycotaxon 16(1): 157-161.

HYPOGYMNIA ENTEROMORPHA TAXONOMY

Pisut, I. 1990. To the distribution of some lichens in central Europe. Bioligia (Bratislava) 45: 685-692.

LICHEN NONVASCULAR CONSERVATION

Pittman, S. 1990. National list of rare lichens. Unpublished report. Not paged.

PILOPHORUS ACICULARIS RAMALINA THRAUSTA TEXOSPORIUM SANCTI-JACOBI XANTHOPARMELIA IDAHOENSIS COLLEMA CURTISPORUM CLADONIA TRANSCENDENS CLADONIA VERRUCULOSA COLLEMA FURFURACEUM HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA LINITA LOBARIA SCROBICULATA PHYSCIA SEMIPINNATA

Rhodes, F. M. [n.d.]. Monitoring air quality in Olympic National Park, Washington, with elemental analysis of lichens. Publisher unknown. Abstract.

AIR QUALITY LICHEN HYPOGYMNIA ENTEROMORPHA

Rice, P. M., R. A. Boldi, C. E. Carlson, P. C. Tourangeau, and C. C. Gordon. 1983. Sensitivity of Pinus ponderosa foliage to airborne phytotoxins: use in biomonitoring. Canadian Journal of Forest Research 13(6): 1083-1091.

AIR QUALITY

Rominger, E. M., and J. L. Oldmeyer. 1989. Early-winter habitat of woodland caribou, Selkirk Mountains, British Columbia. Journal of Wildlife Management 53(1): 238-243.

RANGIFER TARANDUS CARIBOU WOODLAND CARIBOU OLD GROWTH FOREST HABITAT LICHEN WINTERING

Rominger, E. M., and J. L. Oldemeyer. 1990. Early-winter diet of woodland caribou in relation to snow accumulation, Selkirk Mountains, British Columbia, Canada. Canadian Journal of Zoology 68: 2691-2694.

WOODLAND CARIBOU RANGIFER TARANDUS CARIBOU WINTERING FOOD HABITS LICHEN

Rosentreter, R. 1984. The zonation of mosses and lichens along the Salmon River in Idaho. Northwest Science 58(2): 108-117.

IDAHO BRYOPHYTE LICHEN RIPARIAN CLASSIFICATION AQUATIC

Rosentreter, R. 1985. Letter to S. Caicco listing rare plant, lichen, and fungus locations for southern Idaho.

ASCLEPIAS CRYOPTERIS ASTRAGALUS JEJUNUS DIMERESIA HOWELLII ERIOGONUM BREVICAULE STIPA WEBBERI LICHEN FUNGI

Rosentreter, R. 1986. Composition patterns within a rabbit brush (Chrysothamnus) community of the Idaho Snake River Plain. Pages 273-277 in E. D. McArthur and B. L. Welch, compilers, Proceedings- Symposium on the Biology of Artemisia and Chrysothamnus. USDA Forest Service General Technical Report INT-200. Intermountain Research Station, Ogden, UT.

FIRE IDAHO SPECIES BIOLOGY SUCCESSION SHRUBLAND LICHEN

Rosentreter, R. 1987. Collection data for 1987 field season, numbers 4181 through 4405.

MULTI-SPECIES SPECIAL PLANTS LICHEN

Rosentreter, R. 1990. Indicator value of lichen cover on desert shrubs. Pages 282-289 in E. D. McArthur, E. M. Romney, S. D. Smith, and P. T. Tueller, compilers, Proceedings-Symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management. USDA Forest Service General Technical Report INT-276. Intermountain Research Station, Ogden, UT. 351 pp.

LICHEN IDAHO SHRUBLAND STRUCTURE

Rosentreter, R. 1991. Lichens as biological markers. In Proceedings of the Conference: Pesticides in Natural Systems: How Can Their Effects Be Monitored? U. S. Environmental Protection Agency, Region 10. EPA 910/9-91-011.

AIR QUALITY LICHEN

Rosentreter, R., and B. McCune. 1992. Vagrant Dermatocarpon in western North America. The Bryologist 95(1): 15-19.

TAXONOMY SHRUBLAND LICHEN Rosentreter, R. 1992. Vagrant lichens in North America. American Journal of Botany 79(6): 30. Abstract.

LICHEN ASPICILIA FRUTICULOSA XANTHOPARMELIA IDAHOENSIS

Rosentreter, R., and L. Eslick. 1993. Notes on the Bryorias used by flying squirrels for nest construction. Evansia 10(2): 61-63.

LICHEN

Rosentreter, R., L. C. Smithman, and A. DeBolt. 1993. Vernacular lichen names: Swedish names translated to English. Evansia 10(3): 104-111.

LICHEN

Rosentreter, R. 1993. Vagrant lichens in North America. The Bryologist 96(3): 333-338.

ASPICILIA FRUTICULOSA LICHEN DISTRIBUTION XANTHOPARMELIA IDAHOENSIS

Rosentreter, R. 1994. Canopy lichens and how to measure other relevant environmental site conditions in central Idaho. Northwest Science 68(2): 140. Abstract.

LICHEN MONITORING MONTANE FOREST SUBALPINE FOREST

Rosentreter, R. 1994. Lichens of the shrub steppe. Northwest Science 68(2): 141. Abstract.

LICHEN SHRUBLAND

Rosentreter, R. 1994. Lichen diversity in managed forests of the Pacific Northwest. Unpublished report prepared for Eastside Ecosystem Mangement Project. 37+ pp.

LICHEN MONTANE FOREST MANAGEMENT TECHNIQUES NONVASCULAR CONSERVATION STRUCTURE

Rosentreter, R. [n.d.]. New lichen species for Idaho. Unpublished report on file at Idaho Department of Fish and Game, Conservation Data Center, Boise. 7 pp.

LICHEN TEXOSPORIUM SANCTI-JACOBI LOBARIA LINITA LOBARIA SCROBICULATA

Ryan, B. [1995?]. Ecological assessment of lichens of the Columbia Basin of Oregon and Washington. Unpublished report prepared for Eastside Ecosystem Management Project. Paged by section.

LICHEN BIOGEOGRAPHY FUNCTION COMPOSITION STRUCTURE SUCCESSION MONTANE FOREST SUBALPINE FOREST ALPINE GRASSLAND SHRUBLAND COTTONWOOD RIPARIAN ASPICILIA FRUTICULOSA CLADINA RANGIFERINA CLADONIA TRANSCENDENS HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII LOBARIA LINITA PILOPHORUS ACICULARIS PSEUDOCYPHELLARIA ANTHRASPIS TEXOSPORIUM SANCTI-JACOBI

Schaefer, J. A., and W. O. Pruitt, Jr. 1991. Fire and woodland caribou in Southeastern Manitoba. Wildlife Monographs 116: 1-39.

WOODLAND CARIBOU RANGIFER TARANDUS CARIBOU HABITAT FOOD HABITS LICHEN FIRE

Schroeder, G. J., and N. E. Schroeder. 1972. Three Strictaceae previously unreported from Idaho. The Bryologist 75: 101-102.

LOBARIA HALLII PSEUDOCYPHELLARIA ANTHRASPIS DISTRIBUTION IDAHO

Schroeder, G. J., T. L. Esslinger, D. E. Anderegg, and N. E. Schroeder. 1973. Seventy lichen species previously unreported from Idaho. Journal of the Idaho Academy of Science 9: 1-6.

DISTRIBUTION LICHEN

Schroeder, N. E., G. J. Schroeder, and D. E. Anderegg. 1987. Catalog of lichens of Idaho. The Bryologist 78: 32-43.

CHECKLIST IDAHO LICHEN HYPOGYMNIA ENTEROMORPHA LOBARIA HALLII PILOPHORUS ACICULARIS

PSEUDOCYPHELLARIA ANTHRASPIS

Scott, M. D. 1985. The woodland caribou. Pages 494-507 in The Audubon Wildlife Report.

RANGIFER TARANDUS CARIBOU WOODLAND CARIBOU BIOLOGY DISTRIBUTION REPRODUCTION WINTERING HABITAT LICHEN FOOD HABITS TAXONOMY IMPACTS-HUMAN MANAGEMENT TECHNIQUE CONSERVATION BIOLOGY

Sigal, L. L., and T. H. Nash III. Lichen communities on conifers in Southern California mountains: an ecological survey relative to oxidant air pollution. Ecology 64(6): 1343-1354.

AIR QUALITY LICHEN HYPOGYMNIA ENTEROMORPHA MONTANE FOREST

Smith, C. W. 1991. Lichen conservation in Hawaii. Pages 35-45 in D. J. Galloway, editor, Tropical lichens: Their systematics, conservation, and ecology. Systematics Association Special Volume No. 43. Clarendon Press, Oxford.

NONVASCULAR CONSERVATION

Soderstrom, L. 1988. The occurrence of epixylic bryophyte and lichen species in an old natural and a managed forest stand in northeastern Sweden. Biological Conservation 45: 169-178.

MANAGEMENT TECHNIQUES COMMUNITY ECOLOGY NONVASCULAR CONSERVATION

Sorenson, E. R. 1986. Ecology and distribution of ribbed fens in Maine and their relevance to the Critical Areas Program. Unpublished report prepared for the Maine Critical Areas Program, State Planning Office, Augusta. 171 pp.

PEATLAND SCIRPUS CAESPITOSUS CAREX CHORDORRHIZA CAREX LIVIDA DROSERA INTERMEDIA SALIX PEDICELLARIS BRYOPHYTE COMPOSITION STRUCTURE FUNCTION SCIRPUS HUDSONIANUS SCHEUCHZERIA PALUSTRIS VACCINIUM OXYCOCCOS

RHYNCHOSPORA ALBA MUHLENBERGIA RACEMOSA

Spence, J. R. 1985. Checklist of the mosses of Grand Teton National Park and Teton County, Wyoming. Great Basin Naturalist 45(1): 124-126.

CHECKLIST BRYOPHYTE

Spence, J. R. 1986. Bryum calobryoides, a new species from western North America. The Bryologist 89(3): 215-218.

TAXONOMY BRYUM CALOBRYOIDES

Spence, J. R. 1987. The montane moss flora of southwestern British Columbia and northwestern Washington State. Journal of the Hattori Botanical Labratory 62: 417-483.

BRYUM CALOBRYOIDES TAYLORIA SERRATA DISTRIBUTION FLORA

Spence, J. R. 1988. Bryum Hedw. (Bryaceae) in western North America. The Bryologist 91(2): 73-85.

BRYUM CALOBRYOIDES TAXONOMY

Steele, A. 1974. Moss communities of central Idaho forests. Unpublished thesis, University of Idaho, Moscow. 52 pp.

CLASSIFICATION IDAHO BRYOPHYTE SUBALPINE FOREST MONTANE FOREST HELODIUM BLANDOWII

Steele, A. 1978. Bryophyte communities of central Idaho forests. Northwest Science 52(4): 310-322.

CLASSIFICATION STRUCTURE COMPOSITION MONTANE FOREST SUBALPINE FOREST BRYOPHYTE HELODIUM BLANDOWII

Stenroos, S. 1990. Cladonia luteoalba - an enigmatic Cladonia. Karstenia 30: 27-32.

TAXONOMY DISTRIBUTION CLADONIA LUTEOALBA

Steven, G., and D. G. Long. 1989. An update on the status of Buxbaumia aphylla on bings in central Scotland. Transactions of the Botanical Society of Edinburgh 45: 389-395.

BUXBAUMIA APHYLLA STATUS SURVEY DISTRIBUTION

Stolte, K., D. Mangis, R. Doty, and K. Tonnessen, technical coordinators. 1993. Lichens as bioindicators of air quality. USDA Forest Service General Technical Report RM-224. Rocky Mountain Forest and Range Experiment Station, Fort Collins CO. 131 pp.

AIR QUALITY LICHEN

Taylor, J. 1972. The habitat and orientation of capsules of Buxbaumia aphylla in the Douglas Lake region of Michigan. The Michigan Botanist 11: 70-73.

BUXBAUMIA APHYLLA SPECIES BIOLOGY

Thomas, A. E., and R. Rosentreter. 1992. Utilization of lichens by pronghorn antelope in three valleys in east-central Idaho. Bureau of Land Management Technical Bulletin No. 92-3, Idaho State Office, Boise.

ANTILOCAPRA AMERICANA PRONGHORN FOOD HABITS INEL IDAHO LICHEN SHRUBLAND AUTECOLOGY

Thor, G. 1993. Letter outlining the IAL Committee for Conservation of Lichens (ICCL) and questionnaire for information concerning lichens and lichen rich habitats. On file at Idaho Department of Fish and Game, Conservation Data Center, Boise. Not paged.

NONVASCULAR CONSERVATION

Thor, G. 1995. Letter and questionnaire requesting information to establish the IUCN Red List of lichens. On file at Conservation Data Center, Idaho Department of Fish and Game, Boise.

NONVASCULAR CONSERVATION LICHEN

Tonsberg, T. 1980. Contribution to the lichen flora of Majorica (Spain). Norwegian Journal of Botany 28: 193-198.

COLLEMA FURFURACEUM DISTRIBUTION

Toren, D., and L. L. Sigal. 1974. Notes and news: New distribution records of mosses in California. Madrono 22: 311.

BUXBAUMIA PIPERI DISTRIBUTION

Trass, H., and T. Randlane. 1986. Lichen species of Estonia in need of protection. Folia Cryptogamica Estonica 21: 1-3.

NONVASCULAR CONSERVATION LOBARIA SCROBICULATA

Trass, H., and T. Randlane. 1987. Extinct macrolichens of Estonia. Folia Cryptogamica Estonica 25: 1-7.

NONVASCULAR CONSERVATION

U.S. Department of Interior, Bureau of Land Management. 1992. BLM in Idaho - special edition: botany in Idaho. U.S. Department of Interior, Bureau of Land Management, Idaho State Office.

MIRABILIS MACFARLANEI LICHEN PRIMULA ALCALINA

USDA Forest Service and USDI Bureau of Land Management. 1994. Final supplemental environmental impact statement on management of habitat for late-successional and old-growth forest related species within the range of the northern spotted owl. Washington, DC.

NONVASCULAR CONSERVATION RAMALINA THRAUSTA PSEUDOCYPHELLARIA ANTHRASPIS LOBARIA SCROBICULATA LOBARIA HALLII LOBARIA LINITA

Virginia Game Department. 1990. Distribution and status of the northern flying squirrel in Virginia. Annual Report, Virginia Nongame and Endangered Wildlife Investigations. Virginia Department of Game and Inland Fisheries.

NORTHERN FLYING SQUIRREL GLAUCOMYS SABRINUS SOUTHERN FLYING SQUIRREL GLAUCOMYS VOLANS SURVEY METHOD REPRODUCTION LICHEN HABITAT DISTRIBUTION

Vives, J. C. 1974. Additions to the moss flora of Catalonia. Collectanea Botanica "A Barcinonesi Botanico Instituto Edita" 9: 195.

HOOKERIA LUCENS DISTRIBUTION

Welch, W. H. The Hookeriaceae of the United States and Canada. Bryologist 65(1): 1-25.

HOOKERIA LUCENS BRYOPHYTE TAXONOMY DISTRIBUTION

West, S. D. 1973. Buxbaumia aphylla in Alaska. The Bryologist 76: 565-566.

BUXBAUMIA APHYLLA DISTRIBUTION Wheeler, G. A. 1985. Buxbaumia aphylla Hedw. new to Minnesota and notes on some other mosses little known in the state. The Michigan Botanist 24: 77-79.

DISTRIBUTION BUXBAUMIA APHYLLA

Won, S. H. 1976. Annotated checklist of the Hepatics of Idaho. The Bryologist 79: 422-436.

SPHAEROCARPUS HIANS DISTRIBUTION BRYOPHYTE CHECKLIST